PhD course in Probability, Home Assignment 2: Simple Random Walks, Percolation and Fractal Percolation

This home assignment consists of 5 exercises. The deadline is on April 22 2015. On the 23rd the solutions will be posted online, and anything handed in later will give 0 points.

1. SRW: Consider a random walk $(S_n)_{n\geq 1}$ on \mathbb{Z}^d , for which the step size X has distribution

$$\mathbb{P}(X = e_j) = \frac{p}{d}, \ \mathbb{P}(X = -e_j) = \frac{1-p}{d}$$

where $p \in [0, 1]$ (when p = 1/2, this is what we referred to as a simple symmetric random walk). Prove that for any $p \neq 1/2$ and any $d \geq 1$, the random walk $(S_n)_{n\geq 1}$ is transient. Do this by using the combinatorial approach that we used to prove that ssrw is recurrent when d = 1, 2.

2. SRW: Consider two random walks $(S_n)_{n\geq 1}$ and $(T_n)_{n\geq 1}$ on \mathbb{Z}^d , for which the step size X has distribution

$$\mathbb{P}(X=e_j) = \frac{1}{2d}, \ \mathbb{P}(X=-e_j) = \frac{1}{2d}$$

Assume further that the two random walks are independent, and that $S_0 = 0$ and $T_0 = x$ for some $x \in \mathbb{Z}^d$. We say that $(S_n)_{n \ge 1}$ and $(T_n)_{n \ge 1}$ meets, if there exists $n \ge 1$ such that $S_n = T_n$. Prove the following

- (a) If $||x|| := |x_1| + \cdots + |x_d|$ is odd, then for every $d \ge 1$, $\mathbb{P}(S_n = T_n \text{ for some } n \ge 1) = 0$.
- (b) If ||x|| is even, then for every $d \ge 1$, $\mathbb{P}(S_n = T_n \text{ for some } n \ge 1) > 0$. When is this probability 1?
- 3. Percolation and Moment methods: We have that for any random variable X taking values in $\{0, 1, 2, \ldots\}$

$$\mathbb{P}(X > 0) \le \mathbb{E}[X]. \tag{1}$$

Furthermore, by an easy application of Cauchy-Schwarz (you should check this, but its not a part of the assignment)

$$\mathbb{P}(X > 0) \ge \frac{\mathbb{E}[X]^2}{\mathbb{E}[X^2]}.$$
(2)

These elementary upper and lower bounds on $\mathbb{P}(X > 0)$ can be surprisingly useful as we shall see in this exercise.

Consider a tree in which the root o has d children, each of which have d children etc. Denote this tree by \mathbb{T}^d . The case d = 2 is a binary tree which we encountered in the first home assignment. Obviously, \mathbb{T}^d is a deterministic tree, but by performing percolation on it, we create a Galton-Watson tree (which we discussed in the beginning of the course) if we consider the open component of the

root. You can choose to do either edge or site percolation, but they are basically equivalent on trees. For any $x, y \in \mathbb{T}^d$, let $\{x \leftrightarrow y\}$ denote the event that there exists an open path from x to y. Consider the event $\{o \leftrightarrow \infty\}$, which is simply the event that there exists an unbounded open path starting from o.

Let

$$Z_n = |\{x \in \mathbb{T}_n^d : o \leftrightarrow x\}|$$

where \mathbb{T}_n^d is the set of vertices of \mathbb{T}^d at graph distance *n* from the root. Obviously, $\{o \leftrightarrow \infty\} = \bigcap_{n=1}^{\infty} \{Z_n > 0\}.$

- (a) Prove that $\lim_{n\to\infty} \mathbb{P}(Z_n > 0) = 0$, when p < 1/d by using (1).
- (b) Prove that $\lim_{n\to\infty} \mathbb{P}(Z_n > 0) > 0$, when p > 1/d by using (2).

Hint: It can be useful to observe that

$$Z_n = \sum_{x \in \mathbb{T}_n^d} I(o \leftrightarrow x)$$

and to think about what $\mathbb{P}(o \leftrightarrow x, o \leftrightarrow y)$ is.

The above is consistent with what we did at the start of the course, but we can do better! Consider a sequence $(a_n)_{n=0}^{\infty}$ such that $a_n \in \{1, 2, \ldots\}$ for every n, and let $A_n := \prod_{k=0}^{n-1} a_k$. Consider the tree T starting with a root, and let the root have a_0 children who in turn all have a_1 children etc. Using analogous notation to above,

- (c) Prove that $\lim_{n\to\infty} \mathbb{P}(Z_n > 0) = 0$, when $p < 1/\liminf_n A_n^{1/n}$ by using (1).
- (d) Prove that $\lim_{n\to\infty} \mathbb{P}(Z_n > 0) > 0$, when $p > 1/\liminf_n A_n^{1/n}$ by using (2).

Remark: The results (c) and (d) are generalizations of the results in (a) and (b). It is good to do (a) and (b) first to find the right idea for how to prove (c) and (d), but only solving (c) and (d) will of course give full credit.

4. Percolation: Consider percolation on the vertices of the triangular lattice (this is the model that we have studied in class) with density $p < p_c$. Let B_n be the box of side length n centered at the origin (see Figure 1). In order to avoid degenerate situations, we assume that n is odd.

Let $o \leftrightarrow \partial B_n$ denote the event that there exists an open path from the origin to the boundary of the "box" B_n Furthermore, let $\theta_n(p) := \mathbb{P}(o \leftrightarrow \partial B_n)$.

- (a) Prove that there exists a constant $C < \infty$ such that $\theta_n(1/2) \ge 1/(Cn)$ for every (odd) $n \ge 1$.
- (b) Prove that there exist some constants $\nu > 0$ and $C < \infty$ such that for every (odd) $n \ge 1$, and $p \le 1/2$,

$$\theta_n(p) \le C n^{-\nu}.$$

Hints: For (a): draw a picture and compare to the crossing events we used in class. For (b): circuits!

5. Fractals: Consider the fractal percolation model discussed in class, and assume for convenience that d = 2 and N = 3. Recall the notation $\mathcal{C}^1(p) \supset \mathcal{C}^2(p) \supset \cdots$, and as usual define

$$\mathcal{C}(p) := \bigcap_{k=1}^{n} \mathcal{C}^{k}(p).$$



Figur 1: This is the box B_{11} in the notation from class. We also see a configuration of open (black) and closed (white) vertices in this box. The origin is that white guy in the middle.

(a) Consider the set F of "corners", i.e.

$$F := \bigcup_{k=1}^{\infty} \bigcup_{l_1=0}^{3^k} \bigcup_{l_2=0}^{3^k} \left\{ \left(\frac{l_1}{3^k}, \frac{l_2}{3^k} \right) \right\}.$$

The set F consists of all corners of any box on any scale in the fractal construction. Prove that

$$\mathbb{P}(F \cap \mathcal{C}(p) \neq \emptyset) = 0.$$

(b) Consider the lattice $G = (\mathbb{Z}^2, \mathbb{E}^2)$ (i.e. not the hexagonal lattice from class). Perform *site* percolation on G, i.e. pick a random configuration $\omega \in \{0, 1\}^{\mathbb{Z}^2}$ (in the first lecture on percolation, we considered edge percolation on this graph). Let

$$B_n := \mathbb{Z}^2 \cap [0, n]^2.$$

so that B_n is a box of side length n. We let $H^o(B_n)$ denote the event that there exists a left-right open crossing of B_n . Define

$$p_{c,site} := \sup\{p > 0 : \limsup_{n \to \infty} \mathbb{P}_p(H^o(B_n)) = 0\}.$$

Define also

$$HF(p) := \{ \mathcal{C}(p) \text{ contains a L-R crossing of } [0,1]^2 \}$$

(H=Horizontal, F=Fractal), so that with notation from class, $\bar{\varphi}(p) = \mathbb{P}(HF(p))$. Let

$$\bar{p}_c := \inf\{p > 0 : \bar{\varphi}(p) > 0\}.$$

Use part (a), i.e. the fact that with probability one, no corners are in the fractal set, to prove that

 $\bar{p}_c \ge p_{c,site}.$

Hint: Scaling invariance!

Remark: With $\theta_{site}(p)$ defined as in class, but considering site-percolation on \mathbb{Z}^2 , it is known that $p_{c,site}$ is in fact equal to

$$\inf\{p > 0 : \theta_{site}(p) > 0\}.$$

This is a non-trivial statement, which is not needed for the exercise.