

The coinvariant algebra and representation types of blocks of category \mathcal{O} *

Th. Brüstle, S. König and V. Mazorchuk[†]

Abstract

Let \mathfrak{G} be a finite dimensional semisimple Lie algebra over the complex numbers. Let A be the finite dimensional algebra of a (regular or singular) block of the BGG-category \mathcal{O} . By results of Soergel [20], A has a combinatorial description in terms of a subalgebra C_0 of the coinvariant algebra C . In [18], an embedding has been constructed from C_0 -mod into the category $\mathcal{F}(\Delta)$ of A -modules having a Verma flag. This is the main tool for our classification of $\mathcal{F}(\Delta)$ into finite, tame and wild representation type. As a consequence we also obtain a classification of A -mod into finite, tame and wild representation type, thus reproving a recent result of Futorny, Nakano and Pollack [10].

1 Introduction and statement of results

Let \mathfrak{G} be a finite dimensional semisimple Lie algebra over the complex numbers. Fix a triangular decomposition, $\mathfrak{G} = \mathfrak{N}_- \oplus \mathfrak{h} \oplus \mathfrak{N}_+$. Then the BGG-category \mathcal{O} , which has been defined in [4], decomposes into a direct sum of indecomposable subcategories, called *blocks*. By a result of Soergel ([20]) we can restrict our attention from now on to blocks with integral support, since any other block is equivalent to one with integral support (note, however, that applying such an equivalence may involve changing \mathfrak{G} and thus also changing the root system). Each block is equivalent to the module category of a finite dimensional associative algebra A which is unique up to Morita equivalence. The full subcategory of A -mod consisting of modules having a Verma flag is denoted by $\mathcal{F}(\Delta)$. Associated with a block having integral support are the following combinatorial data: an antidominant integral weight λ , the Weyl group W and the stabilizer subgroup W_0 of W fixing λ . Denote the root systems corresponding to W and to W_0 by Φ and Φ_0 , respectively. Our classifications will depend on these root systems only.

*AMS Subject Classification: 16G60, 17B10

[†]Alexander von Humboldt fellow at Bielefeld University

Theorem 1. *Fix a block and its finite dimensional algebra A . Then the representation type of the category $\mathcal{F}(\Delta)$ is given by the following table. All cases not listed are wild.*

| <i>representation type</i> | Φ | Φ_0 |
|----------------------------|-------------------|-----------------|
| <i>finite</i> | <i>any</i> Φ | $\Phi_0 = \Phi$ |
| | A_1 | \emptyset |
| | A_2 | A_1 |
| | A_3 | A_2 |
| | B_2 | A_1 |
| <i>tame</i> | A_4 | A_3 |
| | $A_1 \times A_1$ | \emptyset |
| | A_5 | A_4 |
| | B_3 | B_2 |

Of course, an equality $\Phi = \Phi_0$ occurs precisely for A a simple algebra.

As a consequence of the proof of Theorem 1 we also obtain:

Theorem 2. *Fix a block and its finite dimensional algebra A . Then the representation type of the category $A\text{-mod}$ is given by the following table. All cases not listed are wild.*

| <i>representation type</i> | Φ | Φ_0 |
|----------------------------|-------------------|-----------------|
| <i>finite</i> | <i>any</i> Φ | $\Phi_0 = \Phi$ |
| | A_1 | \emptyset |
| | A_2 | A_1 |
| <i>tame</i> | A_3 | A_2 |
| | B_2 | A_1 |

Theorem 2 has been proved in [10] by completely different methods. The formulation there is for simple Lie algebras. However, our more general formulation follows easily from the results in [10].

Our method is based on Soergel's [20] description of blocks of \mathcal{O} as endomorphism rings over certain subalgebras C_0 of coinvariant algebras and on a result in [18] which states that the module category of C_0 embeds into $\mathcal{F}(\Delta)$ (we reprove this result here for the sake of completeness). The algebra C_0 is local and commutative and explicit computations therein can be easily performed using a basis coming from Schubert calculus. Therefore, the category $C_0\text{-mod}$ is seen to be wild unless we are in one of a small number of limited situations where we really have to study the categories $A\text{-mod}$ and $\mathcal{F}(\Delta)$. These situations have to be dealt with case by case. (A full classification of $C_0\text{-mod}$ into finite, tame and wild type has been obtained by I.Gordon and A.Premet [12].)

In Section 2 we collect some information on coinvariant algebras and their relation to blocks of \mathcal{O} . In Section 3 we consider some special cases, starting with the regular block for $\mathfrak{sl}_2 \times \mathfrak{sl}_2$. In Section 4 a proof of Theorem 1 and of Theorem 2 is given.

2 Coinvariant algebras

We fix \mathfrak{G} , \mathcal{O} , the root system Φ and the Weyl group W as before. The Weyl group W is a Coxeter group with length function l . Moreover, we fix a subset, Θ , of the set of simple roots and denote by W_0 the “parabolic” subgroup of W generated by the corresponding simple reflections. Its root system is called Φ_0 . The Weyl group acts as a reflection group on the complex vector space V and thus on the symmetric algebra $S(V)$. By definition, the *coinvariant algebra* C is the quotient of $S(V)$ modulo the two-sided ideal generated by homogenous W -invariants of degree at least one. By C_0 we denote the (graded) subalgebra of C consisting of W_0 -invariant elements.

It is well-known that C is isomorphic to the cohomology algebra of the flag manifold \mathcal{G}/\mathcal{B} . Therefore, Schubert calculus provides us with a special basis of C , which contains a basis of C_0 as well. This has been worked out in Hiller’s book [14], from which we take the following information (Chapter IV, in particular Sections 3 and 4).

There is a complete set of left coset representatives of W_0 in W consisting of elements w such that each w has minimal length in its coset. These are precisely those w which satisfy $l(ws) = l(w) + 1$ for all simple reflections $s \in \Theta$. The algebra C_0 has a basis consisting of elements X_w indexed by these coset representatives, such that the following properties are satisfied: The element X_w is homogenous of degree l . For a simple reflection, $s = s_\alpha$, and any element, $w \in W$, the product $X_s X_w$ in C is given by the *Pieri formula* (where the ω_α are fundamental weights):

$$X_s X_w = \sum_{\beta \in \Phi^+, l(ws_\beta) = l(w) + 1} (\beta^\vee, \omega_\alpha) X_{ws_\beta}. \quad (1)$$

If s and w are coset representatives, then this is the product formula in C_0 . In particular, if w is chosen in such a way that the product ws is longer than w , then the product $X_s X_w$ is not zero. In fact, in this case X_{ws} occurs with a non-zero coefficient in the right hand term.

Let us now recall the role of C and C_0 in describing blocks of \mathcal{O} . Fix a block by choosing an antidominant integral weight λ . Its stabilizer is a subgroup, W_0 . Then, by Soergel’s *Struktursatz 2* and *Endomorphismensatz 3* from [20], the finite dimensional algebra A associated with this block is the endomorphism ring of a certain module, P , (the “big projective module”) over C_0 :

Theorem 3 (Soergel, [20]). $A \simeq \text{End}_{C_0}(P)$.

We remark that C_0 is a symmetric algebra. In fact, by Soergel’s result it is the endomorphism ring over A of the unique indecomposable projective–injective module P . Hence C_0 is self–injective. As a subalgebra of C , the algebra C_0 is commutative, thus it is symmetric.

Clearly, the module category of C_0 can be embedded into this block. From [18] it follows that even the following is true:

Theorem 4. *The category $C_0\text{-mod}$ embeds into $\mathcal{F}(\Delta)$.*

Proof. This follows by combining Theorem 4 in [18] with Theorem 1 in [18]. Here, we give another - more direct - proof which is built up from arguments already used in [18].

By Soergel's result, the algebra C_0 is the endomorphism ring of the projective-injective A -module P which we can write as Ae for some primitive idempotent $e \in A$. Thus we can identify C_0 with eAe . It is well-known (see e.g. [1]) that the category $eAe\text{-mod}$ is equivalent to the full subcategory of $A\text{-mod}$ consisting of modules M which are P -copresented, i.e. there is an exact sequence $0 \rightarrow M \rightarrow P_1 \xrightarrow{\varphi} P_2$ with $P_1, P_2 \in \text{add}(P)$ direct sums of copies of P . Fix such a module M and a copresentation. It will be enough to show that M has a Verma flag.

Let F be a minimal submodule of P_1 such that it has the following properties: F contains M ; there exists a Δ -filtration $F_0 \subset F_1 \subset \dots \subset P_1$ of P_1 such that F equals some F_i . We are going to show that F equals M .

Let us recall some well-known properties of blocks of \mathcal{O} (see e.g. [6, 4]). The projective module P has a Verma flag. There exists a unique simple Verma module, L , in this block which is the socle of each Verma module. Moreover, L is both the socle and the top of P .

In particular, the image of $\varphi : P_1 \rightarrow P_2$ has a socle which is a direct sum of copies of L . By the minimality of F , the quotient F/M cannot have a composition factor isomorphic to L . Since M is sent to zero by φ , F must be in the kernel as well. This implies equality $F = M$. \square

The embedding preserves isomorphisms and indecomposability. As observed in [18], we therefore have the following lower bound for representation types:

Corollary 1. *If $C_0\text{-mod}$ has infinite representation type, then $\mathcal{F}(\Delta)$ and $A\text{-mod}$ also have infinite type. If $C_0\text{-mod}$ is wild, then $\mathcal{F}(\Delta)$ and $A\text{-mod}$ are wild as well.*

Here we use the following definition of wild representation type for $\mathcal{F}(\Delta)$: The module category $A\text{-mod}$ is wild if there exists an exact functor $F : \mathbb{C}[x, y]\text{-mod} \rightarrow A\text{-mod}$ that preserves non-isomorphism and indecomposability. Now, the exact subcategory $\mathcal{F}(\Delta)$ of $A\text{-mod}$ is called wild if there exists an exact functor F as above that factors through $\mathcal{F}(\Delta)$. Drozd's tame and wild theorem [8] applies to modules having a Verma flag, thus each $\mathcal{F}(\Delta)$ is either wild in the sense defined above or else it is tame.

3 Some explicit examples

In this section we consider some algebras which will appear in special cases where the lower bound from Corollary 1 cannot be used.

3.1 The case $A_1 \times A_1$

If the root system is of type $A_1 \times A_1$, then C_0 has two generators and \mathbb{C} -dimension four. In fact, $C_0 \simeq \mathbb{C}[x, y]/(x^2, y^2)$. Then C_0 is known to be tame [2]; in fact, it is a string algebra ([11, 19]) whose Auslander-Reiten quiver is well-known.

The dimension of C_0 coincides with the number of cosets of W_0 in W . Therefore, this case occurs only for the root system $A_1 \times A_1$ with trivial W_0 . Picking $(\Phi, \Phi_0) = (B_2, A_1)$ yields the same dimension, but in this case C_0 is generated by one element.

The algebra A by definition is a tensor product of two copies of the principal block B of $\mathfrak{sl}(2)$, i.e. of type A_1 . By quiver and relations the last one is given as follows:

$$B: \quad a \bullet \begin{array}{c} \xrightarrow{\alpha} \\ \xleftarrow{\beta} \end{array} \bullet b \quad \begin{array}{l} \text{modulo the ideal generated} \\ \text{by the relation: } \beta \cdot \alpha = 0. \end{array}$$

Therefore, the algebra A has the following quiver and relations:

$$A: \quad \begin{array}{ccc} a \bullet & \begin{array}{c} \xrightarrow{\alpha} \\ \xleftarrow{\beta} \end{array} & \bullet b \\ \gamma \downarrow \uparrow \delta & & \gamma \downarrow \uparrow \delta \\ c \bullet & \begin{array}{c} \xrightarrow{\alpha} \\ \xleftarrow{\beta} \end{array} & \bullet d \end{array} \quad \begin{array}{l} \text{modulo the ideal generated} \\ \text{by the following relations:} \\ \beta \cdot \alpha = 0, \quad \delta \cdot \gamma = 0, \\ \alpha \cdot \gamma = \gamma \cdot \alpha, \quad \alpha \cdot \delta = \delta \cdot \alpha, \\ \beta \cdot \gamma = \gamma \cdot \beta, \quad \beta \cdot \delta = \delta \cdot \beta. \end{array}$$

Here are the composition series of the indecomposable projective modules:

$$P(a) = \begin{array}{ccc} & a & \\ b & & c \\ & d & \end{array}, \quad P(b) = \begin{array}{ccc} & b & \\ d & & a \\ & d & c \end{array}, \quad P(c) = \begin{array}{ccc} & c & \\ b & & d \\ & d & c \end{array}, \quad P(d) = \begin{array}{ccc} & d & \\ b & & c \\ & b & c \\ & d & \end{array}.$$


The projective resolutions of the standard modules look as follows:

$$\begin{aligned} 0 \rightarrow P(a) \rightarrow \Delta(a) = \begin{array}{ccc} & a & \\ b & & c \\ & d & \end{array} \rightarrow 0 \\ 0 \rightarrow P(a) \rightarrow P(b) \rightarrow \Delta(b) = \begin{array}{ccc} & b & \\ d & & a \\ & d & c \end{array} \rightarrow 0 \\ 0 \rightarrow P(a) \rightarrow P(c) \rightarrow \Delta(c) = \begin{array}{ccc} & c & \\ b & & d \\ & d & c \end{array} \rightarrow 0 \\ 0 \rightarrow P(a) \rightarrow P(b) \oplus P(c) \rightarrow P(d) \rightarrow \Delta(d) = \begin{array}{ccc} & d & \\ b & & c \\ & b & c \\ & d & \end{array} \rightarrow 0 \end{aligned}$$

Lemma 1. *The algebra A is wild.*

Proof. The universal covering \tilde{A} of the algebra A is given by a quiver, \tilde{Q} , which is a two-dimensional lattice with arrows labeled as in the figure below, and satisfying the same

relations as for the algebra A . We also marked in the figure a convex subcategory, E , (supported by the 10 thick vertices) of the covering \tilde{A} . As E is contained in Unger's list of wild algebras [21], we conclude that \tilde{A} and therefore also A is wild. \square



We now turn to investigate the category $\mathcal{F}(\Delta)$. Unfortunately, it is sandwiched between the tame category $C_0\text{-mod}$ and the wild module category $A\text{-mod}$, hence we have to use different methods to determine the representation type of $\mathcal{F}(\Delta)$ in this case. We therefore recall from [15] some facts on A_∞ -categories: Let $E = \bigoplus_{i \geq 0} \text{Ext}_A^i(\bigoplus \Delta(x), \bigoplus \Delta(x))$ denote the (graded) Ext-algebra of $\mathcal{F}(\Delta)$. It is endowed with an A_∞ -structure, given by maps $m_n : E^{\otimes n} \rightarrow E$ for $n \geq 2$, where $m_2 : E \otimes E \rightarrow E$ is the Yoneda product. The data $(E, (m_n)_{n \geq 2})$ define a category $\text{filt}(E)$, which we refer to as the category of representations of the A_∞ -algebra E . In case the maps m_n vanish for $n \geq 3$, the algebra (E, m_2) is a graded associative algebra and the category $\text{filt}(E)$ is what is known as the category of representations of the differential graded algebra E (with zero differential, of course).

Proposition 1 (Keller [15], Theorem 7.7). *The category $\mathcal{F}(\Delta)$ is equivalent to $\text{filt}(E)$.*

Thus we can determine the representation type of $\mathcal{F}(\Delta)$ by studying the category $\text{filt}(E)$.

Proposition 2. *The category $\mathcal{F}(\Delta)$ is tame.*

Proof. Of course, we do not verify tameness directly; instead, we reduce this problem to some problem which is known to be tame. First compute the Yoneda product of the Ext-algebra E of $\mathcal{F}(\Delta)$ (e.g. by representing elements in E as shifted morphisms between complexes which are given by the above projective resolutions). It is given by the following (bi-)quiver Q with relations, where the arrows labeled by greek letters describe the degree 0 part of E and the latin letters stand for the degree 1 part.

$$\begin{array}{ccc}
& & \xrightarrow{\alpha_1} \\
\Delta(d) \bullet & \xrightarrow{a_1} & \bullet \Delta(b) \\
\downarrow \beta_1 & \downarrow b_1 & \downarrow \alpha_2 \\
& \searrow c_1 & \\
& c_2 & \\
& \downarrow a_2 & \\
\Delta(c) \bullet & \xrightarrow{b_2} & \bullet \Delta(a) \\
& \xrightarrow{\beta_2} &
\end{array}$$

relations:

$$\begin{aligned}
\alpha_2 \cdot \alpha_1 &= \beta_2 \cdot \beta_1 \\
a_2 \cdot \alpha_1 &= c_1 = \beta_2 \cdot b_1 \\
\alpha_2 \cdot a_1 &= c_2 = b_2 \cdot \beta_1 \\
a_2 \cdot a_1 &= b_2 \cdot b_1.
\end{aligned}$$

Since A is quasi-hereditary, the algebra E is directed. In fact, there are no paths in the quiver Q of length > 2 , hence the maps m_n vanish for $n \geq 3$ (c.f. [15], section 3.4). Thus, the category $\text{filt}(E)$ can be described as follows: A representation $V \in \text{filt}(E)$ is given by linear maps V_x for each latin letter x of Q such that the relation $V_{a_2}V_{a_1} = V_{b_2}V_{b_1}$ holds. In other words, the objects of the category $\text{filt}(E)$ are the representations of the quiver (with relations) given by the latin letters.

The arrows indexed by greek letters come into play when we define morphisms: A morphism in $\text{filt}(E)$ from V to W is given by a family of linear maps

$$H = (H_{\Delta(a)}, H_{\Delta(b)}, H_{\Delta(c)}, H_{\Delta(d)}, H_{\alpha_1}, H_{\alpha_2}, H_{\beta_1}, H_{\beta_2})$$

satisfying $H_{\alpha_2}H_{\alpha_1} = H_{\beta_2}H_{\beta_1}$ such that the following equations hold:

$$\begin{aligned}
H_{\Delta(b)}V_{a_1} &= W_{a_1}H_{\Delta(d)}, & H_{\Delta(a)}V_{a_2} &= W_{a_2}H_{\Delta(b)}, \\
H_{\Delta(c)}V_{b_1} &= W_{b_1}H_{\Delta(d)}, & H_{\Delta(a)}V_{b_2} &= W_{b_2}H_{\Delta(c)}, \\
H_{\beta_2}V_{b_1} + H_{\Delta(a)}V_{c_1} &= W_{c_1}H_{\Delta(d)} + W_{a_2}H_{\alpha_1}, \\
H_{\alpha_2}V_{a_1} + H_{\Delta(a)}V_{c_2} &= W_{c_2}H_{\Delta(d)} + W_{b_2}H_{\beta_1}.
\end{aligned}$$

In order to show finally that $\text{filt}(E)$ is tame, we use the classical reduction method for differential graded categories ([16, 17]): Consider the (graded) subalgebra G of E generated by the arrows c_1 and c_2 . The reduction of E with respect to G leads to a new differential graded category, K , whose category of representations is equivalent to $\text{filt}(E)$.

In our case, the quotient E/G is given by a commutative square, a well-known representation-finite algebra. It is therefore easy to compute the category K and it turns out that K is given by a bundle of chains in the sense of [3], namely by the following two pairs of posets

$$(x \rightarrow \bullet, \quad y \rightarrow \bullet) \quad \text{and} \quad (\tilde{x} \rightarrow \bullet, \quad \tilde{y} \rightarrow \bullet)$$

with involution $x \mapsto \tilde{x}$ and $y \mapsto \tilde{y}$. Thus, by [3] or equivalently [5], the category of representations of K is tame and its indecomposable objects have been completely classified. \square

3.2 An enlargement of $A_1 \times A_1$

When proving the theorems we have to consider some enlargements of the case $A_1 \times A_1$. We therefore collect in the next two subsections auxiliary statements about quasi-hereditary algebras that are given by composition series of projective modules and by projective resolutions of standard modules. In the proofs later on, quotients and subalgebras of blocks of category \mathcal{O} will appear which have precisely these properties (this in particular will ensure the existence of these algebras). It will turn out that for proving tameness or wildness of the corresponding categories $\mathcal{F}(\Delta)$ the partial information we have on these algebras is sufficient.

The application we are aiming at allows us to make some assumptions which simplify notation. In particular, we only consider algebras with a duality. In category \mathcal{O} terminology the algebras appearing in this context are multiplicity free. Thus all indecomposable projective modules are contained in a unique 'big' projective module. The quotients or subalgebras to be studied now inherit this property which explains another assumption made below.

Thus, let A be a quasi-hereditary algebra (admitting a duality fixing simples) which has five primitive idempotents $\{a, b, c, d, e\}$ such that the standard modules have the following projective resolutions:

$$\begin{aligned}
 0 \rightarrow P(a) \rightarrow \Delta(a) &= \begin{matrix} a \\ b & c \\ d \\ e \end{matrix} \rightarrow 0 \\
 0 \rightarrow P(a) \rightarrow P(b) \rightarrow \Delta(b) &= \begin{matrix} b \\ d \\ e \end{matrix} \rightarrow 0 \\
 0 \rightarrow P(a) \rightarrow P(c) \rightarrow \Delta(c) &= \begin{matrix} c \\ d \\ e \end{matrix} \rightarrow 0 \\
 0 \rightarrow P(a) \rightarrow P(b) \oplus P(c) \rightarrow P(d) \rightarrow \Delta(d) &= \begin{matrix} d \\ e \end{matrix} \rightarrow 0 \\
 0 \rightarrow P(d) \rightarrow P(e) \rightarrow \Delta(e) &= e \rightarrow 0
 \end{aligned}$$

The composition series of the indecomposable projective modules are:

$$P(a) = \begin{matrix} a \\ b & c \\ d \\ e \end{matrix}, \quad P(b) = \begin{matrix} b \\ d & a \\ e & c \end{matrix}, \quad P(c) = \begin{matrix} c \\ a & d \\ b & e \\ d \\ e \end{matrix}, \quad P(d) = \begin{matrix} d \\ b & e & c \\ e & a & d \\ & b & c & e \\ & & d \\ & & e \end{matrix},$$

$$\begin{array}{ccccc}
& & e & & \\
& & d & & \\
& b & e & c & \\
P(e) = & d & a & d & \\
& e & b & c & e \\
& & d & & \\
& & e & &
\end{array}$$

We further suppose that the projectives admit inclusions $P(a) \subset P(b) \subset P(d)$, $P(a) \subset P(c) \subset P(d)$ and $P(d) \subset P(e)$.

Proposition 3. *The category $\mathcal{F}(\Delta)$ is wild.*

Proof. As in the proof of Proposition 2, we consider $E = \bigoplus_{i \geq 0} \text{Ext}_A^i(\bigoplus \Delta(x), \bigoplus \Delta(x))$, the Ext-algebra of the standard modules. Let F be the (graded) subalgebra of E generated by extensions of $\Delta(a) \oplus \Delta(d) \oplus \Delta(e)$, i.e.

$$F = \bigoplus_{i \geq 0} \text{Ext}_A^i(\Delta(a) \oplus \Delta(d) \oplus \Delta(e), \Delta(a) \oplus \Delta(d) \oplus \Delta(e)).$$

We know that E is endowed with an A_∞ -structure such that $\mathcal{F}(\Delta)$ and $\text{filt}(E)$ are equivalent. Our strategy is the following: we show that the A_∞ -structure restricted to F is just given by the Yoneda-product (i.e. $m_n : E^{\otimes n} \rightarrow E$ vanishes for $n \geq 3$ when restricted to $F^{\otimes n}$). Thus, $\text{filt}(F)$ can be viewed as the category of representations of the (differential) graded algebra F and it is a full subcategory of $\text{filt}(E)$. Therefore it suffices to show that $\text{filt}(F)$ is wild. Note that this proof cannot be obtained on the level of quasi-hereditary algebras: the category $\text{filt}(F)$ consists of those objects in $\mathcal{F}(\Delta)$ whose multiplicity of $\Delta(b)$ and $\Delta(c)$ is zero. The corresponding algebra fAf with $f = a + d + e$ however is not quasi-hereditary.

From the projective resolutions of the standard modules and the given embeddings of the projective modules we can easily compute the dimensions of morphism and extension spaces between standard modules. In particular, the Ext^1 -quiver of the graded algebra F looks as follows (here the arrows correspond to basis vectors of the Ext^1 -spaces between the various standard modules):

$$\begin{array}{ccccc}
\bullet & \xrightarrow{x} & \bullet & \xrightarrow{\quad} & \bullet \\
\Delta(e) & & \Delta(d) & & \Delta(a)
\end{array}$$

Since A is quasi-hereditary, the (differential) graded algebra F is directed. Moreover, F admits only three primitive idempotent elements, namely the identity morphisms of $\Delta(e)$, $\Delta(d)$ and $\Delta(a)$. Therefore, by [15, section 3.4], the maps $m_n : E^{\otimes n} \rightarrow E$ vanish when restricted to $F^{\otimes n}$ for $n \geq 3$.

The Ext^1 -quiver of F contains the following wild subquiver Γ :

$$\begin{array}{ccccc}
\bullet & \longrightarrow & \bullet & \xrightarrow{\quad} & \bullet \\
\Delta(e) & & \Delta(d) & & \Delta(a)
\end{array}$$

One easily sees that $\text{Ext}^2(\Delta(a), \Delta(e)) = 0$. Moreover, the standard modules have only trivial endomorphisms, and all morphisms between different standard modules induce morphisms in $\text{filt}(F)$ that involve the arrow x which is not contained in Γ . Therefore the obvious functor $\text{rep } \Gamma \rightarrow \text{filt}(F)$ preserves non-isomorphism and indecomposability. We conclude that the category $\text{filt}(F)$ and hence $\mathcal{F}(\Delta)$ is wild. \square

3.3 Another enlargement of $A_1 \times A_1$

We consider finally a second enlargement of the case $A_1 \times A_1$: Let A be a quasi-hereditary algebra (admitting a duality fixing simples) with primitive idempotents $\{a, a', b, c, d\}$ whose standard modules have the following projective resolutions:

$$\begin{aligned} 0 \rightarrow P(a) \rightarrow \Delta(a) = \begin{matrix} a \\ b & c \\ d \end{matrix} \rightarrow 0 \\ 0 \rightarrow P(a') \rightarrow \Delta(a') = \begin{matrix} a' \\ b & c \\ d \end{matrix} \rightarrow 0 \\ 0 \rightarrow P(a) \oplus P(a') \rightarrow P(b) \rightarrow \Delta(b) = \begin{matrix} b \\ d \end{matrix} \rightarrow 0 \\ 0 \rightarrow P(a) \oplus P(a') \rightarrow P(c) \rightarrow \Delta(c) = \begin{matrix} c \\ d \end{matrix} \rightarrow 0 \\ 0 \rightarrow P(a) \oplus P(a') \rightarrow P(b) \oplus P(c) \rightarrow P(d) \rightarrow \Delta(d) = d \rightarrow 0 \end{aligned}$$

The composition series of the indecomposable projective modules are:

$$P(b) = \begin{matrix} & b & & & \\ d & a & a' & & \\ b & c & c & b & \\ & d & & d & \end{matrix}, \quad P(c) = \begin{matrix} & & c & & \\ a & a' & d & & \\ b & c & c & b & \\ & d & & d & \end{matrix}, \quad P(d) = \begin{matrix} & & & d & \\ & & b & c & \\ d & a & a' & d & \\ & b & c & c & b \\ & d & & d & \end{matrix}.$$

Proposition 4. *The category $\mathcal{F}(\Delta)$ is wild in this case.*

Proof. This works completely analogous to the case we solved before: From the projective resolutions of the standard modules we obtain the following wild subquiver Γ of the quiver of E with all arrows of degree 1:

$$\begin{array}{ccccc} \bullet & \xleftrightarrow{\quad} & \bullet & \xleftrightarrow{\quad} & \bullet \\ \Delta(a) & & \Delta(d) & & \Delta(a') \end{array}$$

It is also clear that $\text{Hom}_A(\Delta(a), \Delta(a'))$ and $\text{Hom}_A(\Delta(a'), \Delta(a))$ are zero, thus the wildness of the subquiver Γ implies that $\mathcal{F}(\Delta)$ is wild. \square

4 Proof of Theorems

We fix A and C_0 as before. Recall that C_0 is a graded symmetric algebra. We will go through several cases depending on Φ_0 . In each case, the first step of the proof consists of determining the representation type of C_0 . If C_0 is wild, then this case is finished. Only few situations remain where C_0 is not wild. In a second step we have to deal with these situations.

In order to decide about the representation type of C_0 we will use the following well-known lemma.

Lemma 2. *A local algebra is wild if its minimal number of generators is at least three.*

Proof. This follows from [13]; see also [9], I.10.10 (a). \square

Now let us distinguish four cases according to the number of simple reflections not contained in Θ , i.e. the corank of Φ_0 in Φ . If this number is zero, then $W = W_0$ and the algebra A is simple. If the corank is three or bigger, then the minimal number of generators of C_0 is at least three, hence by Lemma 2 the algebra C_0 is wild. Thus it remains to consider the two cases of the corank being one or two.

Let us first suppose that Φ_0 has corank two. There are two simple reflections, s_α and s_β , outside of Θ . Denote by w the product $s_\alpha s_\beta$. The four elements $1, s_\alpha, s_\beta$ and w are minimal length representatives of four different cosets of W_0 in W , and each of them defines a basis element in C_0 . If the dimension of C_0 is four, then we are in the case $A_1 \times A_1$ which has been discussed in subsection 3.1. Otherwise the dimension of C_0 must be at least six since it is symmetric. Passing to quotients or subalgebras of A if necessary we arrive at an algebra A as in subsection 3.3 or at an algebra A' which differs from this A by having some multiplicities (of simple modules in Verma modules) bigger than one, i.e. by some arrows occurring with multiplicities. Clearly, the case of A' also falls into wild type. This automatically implies that the original algebra we started with is wild as well. Note that in this case our lower bound may be too weak, since C_0 may be tame (e.g. in type A_2 , see [12]).

It remains to consider the case of Φ_0 having corank one. Denote by s the simple reflection which is not in Θ . In degree one, the graded algebra C_0 consists of the scalar multiples of X_s . If all other degrees are one-dimensional as well, then C_0 is isomorphic to $\mathbb{C}[x]/(x^m)$ for some m . It is obviously of finite representation type, thus our lower bound is trivial.

In this case, we can determine A itself. In fact, by Soergel's double centralizer property (Theorem 3), the algebra A then is the Auslander algebra of C_0 , i.e. the endomorphism ring of the sum of all indecomposable C_0 -modules. By [7], Section 7, the category $\mathcal{F}(\Delta)$ has finite representation type if and only if $m \leq 5$; it is tame for $m = 6$ and wild otherwise. The full category A -mod is of finite type for $m \leq 3$ only, tame for $m = 4$ and wild otherwise.

The number m is the \mathbb{C} -dimension of C_0 , hence it equals the number of cosets of W_0 in W . Consequently, we have the following choices for the root systems (Φ, Φ_0) : For $m = 1$, of course, $\Phi = \Phi_0$ and no further restriction. For $m = 2$ we have (A_1, \emptyset) . For $m = 3$ we have (A_2, A_1) ; $m = 4$ means (A_3, A_2) or (B_2, A_1) ; $m = 5$ occurs for (A_4, A_3) and $m = 6$ happens precisely for (A_5, A_4) and for (B_3, B_2) .

From now on, we may assume that we are not in the situation of Lemma 2, that Φ_0 has corank one and that some graded pieces of C_0 have \mathbb{C} -dimension bigger than one. Denote the smallest degree where this happens by $l + 1$. Note that the dimension of C_0 must be bigger than four in this case. As before, we have a generator X_s associated with a simple reflection, $s \notin \Theta$. If the dimension in degree $l + 1$ is bigger than two, then we are in the situation of Lemma 2. Thus we may assume that the dimension is two. Denote the two standard basis elements by X_{w_1} and X_{w_2} .

If w_1s is longer than w_1 , then the Pieri formula ((1) in Section 2) implies that the product $X_{w_1}X_s$ is not zero. If also w_2s is longer than w_2 , then one of the elements X_{w_1} and X_{w_2} is an additional generator and (passing to quotients or subalgebras if necessary and up to forgetting higher multiplicities) we are in the situation of subsection 3.2.

Suppose now that w_1s is longer than w_1 , but w_2s is shorter than w_2 . Then the element w_2s must be the minimal length representative of its W_0 -coset (if not, write $w_2s = w_0v$ for some $w_0 \in W_0$ and v of length at most $l - 1$; then $w_2 = w_0vs$ cannot be minimal, since its length $l + 1$ is bigger than that of vs). Therefore X_{w_2s} generates the degree l -piece of C_0 . The Pieri formula now tells us that X_{w_2} occurs with a non-zero coefficient in the expansion of the product $X_{w_2s}X_s$. It follows that X_{w_1} is an additional generator and we are again in the situation of subsection 3.2.

Finally, it cannot happen that both w_1s and w_2s are shorter than w_1 and w_2 . If so, by the same arguments as above, the basis elements X_{w_1s} and X_{w_2s} would be linearly dependent. Hence w_1s and w_2s would have to coincide, a contradiction.

This finishes the proofs of Theorems 1 and 2.

Acknowledgements

The research was done during the visit of the third author to Bielefeld University as an Alexander von Humboldt fellow. The financial support of Humboldt Foundation and the hospitality of Bielefeld University are gratefully acknowledged. We are indebted to I. Gordon for pointing out some mistakes in an earlier version.

References

- [1] *M. Auslander*, Representation theory of Artin algebras I. Commun. in Alg. 1 (1974), 177-268.
- [2] *V.A. Bašev*, Representations of the group $Z_2 \times Z_2$ in a field of characteristic 2. (Russian) Dokl. Akad. Nauk SSSR 141 (1961), 1015-1018.

- [3] *V.M.Bondarenko*, Representations of bundles of semichained sets and their applications. (Russian) Algebra i Analiz 3 (1991), no. 5, 38–61; translation in St. Petersburg Math. J. 3 (1992), no. 5, 973–996
- [4] *I.N.Bernstein, I.M.Gelfand and S.I.Gelfand*, A certain category of \mathfrak{G} -modules. (Russian) Funkcional. Anal. i Priložen. 10 (1976), no. 2, 1–8.
- [5] *W.W.Crawley-Boevey*, Functorial filtrations. II. Clans and the Gelfand problem. J. London Math. Soc. (2) 40 (1989), no. 1, 9–30.
- [6] *J.Dixmier*, Enveloping algebras. Revised reprint of the 1977 translation. Graduate Studies in Mathematics, 11. American Mathematical Society, Providence, RI (1996).
- [7] *V.Dlab and C.M.Ringel*, The module theoretical approach to quasi-hereditary algebras. In: H.Tachikawa und S.Brenner (Eds.): Representations of algebras und related topics, London Mathematical Society LN Series 168, 200–224 (1992).
- [8] *Ju.A.Drozd*, Tame and wild matrix problems. (Russian) Representations and quadratic forms (Russian), pp. 39–74, 154, Akad. Nauk Ukrain. SSR, Inst. Mat., Kiev, 1979.
- [9] *K. Erdmann*, Blocks of tame representation type and related algebras. Springer LN 1428 (1990).
- [10] *V. Futorny, D. Nakano and D. Pollack*, Representation type of the blocks of category \mathcal{O} . Preprint (1999).
- [11] *I. M. Gelfand and V. A. Ponomarev*, Indecomposable representations of the Lorentz group. (Russian) Uspehi Mat. Nauk 23 (1968), no. 2 (140), 3–60.
- [12] *I. Gordon and A. Premet*, Block representation type of reduced enveloping algebras. Preprint (2000).
- [13] *A.Heller and I.Reiner*, Indecomposable representations. Illinois J. Math. 5 (1961), 314–323.
- [14] *H. Hiller*, Geometry of Coxeter groups. Pitman Research Notes in Mathematics 54 (1982).
- [15] *B. Keller*, Introduction to A-infinity algebras and modules. Preprint (1999), available via www at “<http://www.math.jussieu.fr/~keller/recent.html>”.
- [16] *A.V.Roĭter and M.M.Kleiner*, Representations of differential graded categories. Representations of algebras (Proc. Internat. Conf., Carleton Univ., Ottawa, Ont., 1974), pp. 316–339. Lecture Notes in Math., Vol. 488, Springer, Berlin (1975).
- [17] *M.M.Kleĭner and A.V.Roĭter*, Representations of differential graded categories. (Russian) Matrix problems (Russian), pp. 5–70. Akad. Nauk Ukrain. SSR Inst. Mat., Kiev (1977).
- [18] *S.König and V.Mazorchuk*, Enright’s completions and injectively copresented modules. Preprint (1999), available via www at “<http://www.mathematik.uni-bielefeld.de/~koenig/>”.
- [19] *C.M.Ringel*, The indecomposable representations of the dihedral 2-groups. Math. Ann. 214 (1975), 19–34.
- [20] *W. Soergel*, Kategorie \mathcal{O} , perverse Garben und Moduln über den Koinvarianten zur Weylgruppe. (in German) [Category \mathcal{O} , perverse sheaves and modules over the coinvariants for the Weyl group], J. Amer. Math. Soc. 3 (1990), no. 2, 421–445.
- [21] *L. Unger*, The concealed algebras of the minimal wild, hereditary algebras. Bayreuther Math. Schr., 31 (1990), 145–154.

Thomas Brüstle, Fakultät für Mathematik, Universität Bielefeld, Postfach 100131, D-33501, Bielefeld, FRG, e-mail: bruestle@mathematik.uni-bielefeld.de

Steffen König, Fakultät für Mathematik, Universität Bielefeld, Postfach 100131, D-33501, Bielefeld, FRG, e-mail: koenig@mathematik.uni-bielefeld.de

Present address: Department of Mathematics and Computer Science, University of Leicester, Leicester LE1 7RH , UK, e-mail: `sck5@mcs.le.ac.uk`

Volodymyr Mazorchuk, Algebra, Mechanics and Mathematics Department, Kyiv Taras Shevchenko University, 64 Volodymyrska st., 01033, Kyiv, Ukraine, e-mail: `mazor@mechmat.univ.kiev.ua`;
`mazor@mathematik.uni-bielefeld.de`

Present address: Department of Mathematics, Chalmers University of Technology and University of Goeteborg, SE-412 96, Sweden, e-mail: `volody@math.chalmers.se`