

FIRST-PASSAGE PERCOLATION WITH EXPONENTIAL TIMES ON A LADDER*

Henrik Renlund[†]

Abstract

We consider first-passage percolation on a ladder, i.e. the graph $\mathbb{N} \times \{0, 1\}$ where nodes at distance 1 are joined by an edge, and the times are exponentially i.i.d. with mean 1. We find an appropriate Markov chain to calculate an explicit expression for the *time constant* whose numerical value is ≈ 0.6827 . This time constant is the long-term average inverse speed of the process. We also calculate the average residual time.

1 Introduction

Consider a graph G with vertex (node) set V and edge set $E \subset V \times V$. An (undirected) edge $e = \langle v, v' \rangle = \langle v', v \rangle$ joins vertex v and v' . A path $\pi(v, v')$ between v and v' , if it exists, is an alternating sequence of vertexes and edges $(v_0, e_1, v_1, \dots, e_n, v_n)$ such that $e_i = \langle v_{i-1}, v_i \rangle$ for $i = 1, \dots, n$, $v_0 = v$ and $v_n = v'$.

Associate with each edge e a non-negative random variable ξ_e . Let the time of a path $\pi(v, v')$ be

$$T\pi(v, v') = \sum_{e \in \pi(v, v')} \xi_e \quad \text{and let} \quad T(v, v') = \inf_{\pi(v, v')} T\pi(v, v')$$

be the shortest time of any path between v and v' . This may be called the first passage time of v' (with respect to v) and is the subject of investigation in first passage percolation, see e.g. [SW78].

We will think of this as a model for a contagious disease and refer to the first passage time of a node as the time of infection. Usually, we think of some node v (or more generally a set of nodes) as infected at time zero. The times $T(w) = T(v, w)$, $w \in V$, lets us know when nodes $w \in V$ are infected.

One natural question is; how fast is the spread of the infection? The present paper considers the equivalent question of the inverse speed of percolation on a ladder, i.e. how long does it take for the infection to spread one more step up the ladder, averaged over time. We also calculate the average inverse speed in another sense; namely the average residual time.

*MSC Primary 82B43. MSC Secondary 60K35.

[†]Math. Department, Uppsala University, P.O. Box 480, 751 06 Uppsala, Sweden

2 First-passage percolation on the ladder

First-passage percolation on a ladder has previously been studied in [FGS06], which gives a method of calculating the time constant when the times associated with edges have a discrete distribution, as well as a method for getting arbitrarily good bounds for the same quantity when the distribution is continuous (and well behaved). Our paper gives an exact formula for the time constant only when the times are exponential and the method we use relies heavily on the special properties of this distribution.

It has been brought to our attention that during the referee period of the present paper [Sch09] has emerged, which also contains the percolation rate (inversion of the time constant), although it is achieved through other methods.

Let L be a ladder, by which we mean a graph with vertex set $V = \mathbb{N} \times \{0, 1\}$ and where the edge set E consists of $\langle v, v' \rangle$ such that v and v' are at distance one from each other, see Figure 1. By \mathbb{N} we mean the set of non-negative integers $\{0, 1, 2, \dots\}$.




Figure 1: Part of the ladder L .

We will consider the model where the nodes $(0, 0)$ and $(0, 1)$ are infected at time zero and each edge is associated with a random variable that has an exponential distribution with mean 1. Random variables associated to different edges are independent.

For a node $v = (x, y)$ we will say that v is at height $x \in \mathbb{N}$ and level $y \in \{0, 1\}$. Recall that $T[(x, y)]$ is the infection time of node (x, y) . Let

$$N_t^{(i)} = \sup\{x : T[(x, i)] \leq t\}, \quad i = 0, 1,$$

i.e. let $N_t^{(0)}$ and $N_t^{(1)}$ denote the height of the infection at level 0 and 1, respectively, at time t .

We will let $N_t = \max\{N_t^{(0)}, N_t^{(1)}\}$ denote the height of the infection, i.e. the largest height of any infected node, at time t .

We aim to calculate the time constant τ , defined as the limiting inverse percolation rate, $1/\tau = \lim_t N_t/t$. We will not do this directly but calculate the percolation rate at “time infinity”, or rather when a process related to N_t has reached stationarity.

More explicitly, we will employ an observation by Alm and Janson [AJ] that arose in connection with their joint work on one-dimensional lattices [AJ90], namely; it might be fruitful to consider the *front* $F_t = |N_t^{(0)} - N_t^{(1)}|$ at time $t \geq 0$. Front thus means the absolute value of the difference in height between the highest infected nodes on each level, see Figure 2.

Now, the process $\{F_t, t \geq 0\}$ behaves like a continuous time Markov chain on \mathbb{N} , with $F_0 = 0$. Suppose that this Markov chain tends to a stationary distribution $\Pi = (\pi_0, \pi_1, \pi_2, \dots)$ on \mathbb{N} . Then at a late time t the process F will be in state 0 with probability π_0 . From this state there are two possible nodes that may be the next infected ones that result in an increase of the N -process, i.e. the intensity is 2 towards a state that increases the height of the infection, since these two nodes are associated with two distinct edges (and thus two distinct exponentially distributed mean 1 random variables). With probability $1 - \pi_0$ the F process is in some other state ($F \geq 1$) from where there is intensity 1 towards a state that increases the height of the infection. Thus, knowing the stationary distribution (or rather knowing π_0) gives us the percolation rate at a late time as

$$2\pi_0 + 1(1 - \pi_0) = 1 + \pi_0. \tag{1}$$

Below we calculate this stationary distribution Π . To do so, it turns out that we may express any π_n in terms of π_0 in the form $\pi_n = a_n\pi_0 - b_n$ where both sequences of (positive) coefficients, a_n and b_n , satisfy a certain recursion (Claim 1) and can be expressed in terms of the Bessel functions of the first and second kind. We stumbled upon the solution, and may not have solved it otherwise, when it was noted that the sequence b_n transformed into $B_n = (b_n - b_{n-1})/n$ is part of sequence A058797, as listed in [Sl07], which is related to the Bessel functions. This relation to Bessel functions was a surprise and we see no reason to expect it in this context. This method can be generalized to percolation on a more general “ladder-like” graph, see [PR]. The sequence b_n is also a subsequence of A056921.




Figure 2: Infected nodes at time t marked as black. Here $N_t = 6$ and $F_t = 2$.

Now, we start by looking at the intensity matrix Q of the process F_t . Consider the case in Figure 2, where $F_t = 2$. Now, there are two edges leading to the node $(5, 1)$ and if this is the next infected, the process will end up in state $F = 1$, i.e. the intensity to this state is 2. There is one edge

each to node $(6, 1)$ and $(7, 0)$ and if these are the next ones infected, the process will end up in state $F = 0$ or $F = 3$, respectively. This argument gives us the third row of the intensity matrix Q . We omit the rest of the details, but it is not hard to see that

$$Q = \begin{pmatrix} -2 & 2 & 0 & 0 & 0 & & & & & & \\ 2 & -3 & 1 & 0 & 0 & & & & & & \\ 1 & 2 & -4 & 1 & 0 & \dots & & & & & \\ 1 & 1 & 2 & -5 & 1 & & & & & & \\ 1 & 1 & 1 & 2 & -6 & & & & & & \\ 1 & 1 & 1 & 1 & 2 & & & & & & \\ & & & \vdots & & & & & & \ddots & \end{pmatrix}. \quad (2)$$

$\{F_t, t \geq 0\}$ is irreducible, so if we can find a distribution $\Pi = (\pi_0, \pi_1, \pi_2, \dots)$ on \mathbb{N} such that $\Pi Q = 0$ then Π is the unique stationary distribution for $\{F_t, t \geq 0\}$, by Markov chain theory. For calculations we henceforth assume that such a distribution Π exists.

Using the equations arising from columns one to three of Q , the condition $\Pi Q = 0$ and $\sum \pi_j = 1$ we get

$$\pi_1 = 3\pi_0 - 1, \quad \pi_2 = 11\pi_0 - 5, \quad \text{and} \quad \pi_3 = 56\pi_0 - 26. \quad (3)$$

Also, taking differences of equations arising from adjacent columns of Q , from column three and upwards, gives the equations

$$0 = \pi_{n-3} - (n+1)\pi_{n-2} + (n+3)\pi_{n-1} - \pi_n, \quad n \geq 4. \quad (4)$$

We set $\pi_n = a_n\pi_0 - b_n$, where $a_1, a_2, a_3, b_1, b_2, b_3$ are defined by (3) for $n = 1, 2, 3$ and, for $n \geq 4$ by (4) with a_n or b_n in place of π_n . Table 1 gives the first nine numbers in the sequences $\{a_n\}$ and $\{b_n\}$.

n	1	2	3	4	5	6	7	8	9
a_n	3	11	56	340	2395	19231	173490	1737706	19136803
b_n	1	5	26	158	1113	8937	80624	807544	8893225

Table 1: The beginning of the sequences $\{a_n\}$ and $\{b_n\}$.

Claim 1. *The sequences $\{a_n\}$ and $\{b_n\}$ both satisfy the recursion*

$$c_n = \frac{c_{n+1} - c_n}{n+1} - \frac{c_n - c_{n-1}}{n}, \quad n \geq 2. \quad (5)$$

Proof. We will use induction. First, check initial conditions, $a_2 = 11$ and $(56 - 11)/3 - (11 - 3)/2 = 11$, $b_2 = 5$ and $(26 - 5)/3 - (5 - 1)/2 = 5$, so equation (5) holds for $n = 2$. Rearranging (5) gives

$$c_{n+1} - c_n = (n+1) \left(c_n + \frac{1}{n} [c_n - c_{n-1}] \right), \quad n \geq 2, \quad (6)$$

and this relation is what we will prove. So assume (6) holds for some $n \geq 2$. We aim to show that it also holds for $n + 1$. So, as $n + 2 \geq 4$, we can use (4). Adding $\pi_n - \pi_{n-1}$ to both sides of, and substituting $n + 2$ for n in (4) gives

$$\begin{aligned} c_{n+2} - c_{n+1} &= c_{n-1} - (n+3)c_n + (n+5)c_{n+1} - c_{n+1} \\ &= (n+2) \left(c_{n+1} + \frac{1}{n+2} [2c_{n+1} - (n+2)c_n - \underbrace{(c_n - c_{n-1})}_{(*)}] \right). \end{aligned} \quad (7)$$

From (6), and the induction hypothesis, we get $c_n - c_{n-1} = (nc_{n+1} - n(n+2)c_n)/(n+1)$, which we apply on (*) above, and

$$\begin{aligned} (7) &= (n+2) \left(c_{n+1} + \frac{1}{n+2} \left[\left(2 - \frac{n}{n+1}\right) c_{n+1} - \left(n+2 - \frac{n(n+2)}{n+1}\right) c_n \right] \right) \\ &= (n+2) \left(c_{n+1} + \frac{1}{n+1} [c_{n+1} - c_n] \right). \end{aligned}$$

□

Now set $C_n = (c_n - c_{n-1})/n$. Then, if c_n satisfies (5), i.e. $c_n = C_{n+1} - C_n$, this implies both

$$C_{n+1} = c_n + C_n \quad (8)$$

and $C_n = C_{n+1} - c_n$. Substitute $n - 1$ for n in the last equation to get

$$C_{n-1} = C_n - c_{n-1}. \quad (9)$$

Then adding (8) and (9) and replacing $c_n - c_{n-1}$ with nC_n gives us the following.

Claim 2. $A_n = (a_n - a_{n-1})/n$ and $B_n = (b_n - b_{n-1})/n$ both satisfy the recursion

$$C_{n+1} + C_{n-1} = (n+2)C_n, \quad n \geq 3. \quad (10)$$

2.1 Bessel functions

The functions

$$J_n(x) = \sum_{k=0}^{\infty} \frac{(-1)^k}{k!(n+k)!} \left(\frac{x}{2}\right)^{n+2k}$$

and

$$\begin{aligned} Y_n(x) &= \frac{1}{\pi} \left[2 \left(\gamma + \ln \frac{x}{2} \right) J_n(x) - \sum_{k=0}^{n-1} \frac{(n-k-1)!}{k!} \left(\frac{x}{2}\right)^{2k-n} \right. \\ &\quad \left. - \sum_{k=0}^{\infty} (\varsigma_k + \varsigma_{k+n}) \frac{(-1)^k}{k!(n+k)!} \left(\frac{x}{2}\right)^2 \right], \end{aligned}$$

where $n \in \mathbb{N}$, $\varsigma_m = \sum_{j=1}^m \frac{1}{j}$ and $\gamma = 0.577\dots$ is Euler's constant, are known as Bessel functions of first and second kind. They both satisfy the recursion

$$C_{n+1}(x) + C_{n-1}(x) = \frac{2n}{x}C_n(x). \quad (11)$$

Define the function

$$\Upsilon(n, m) = \pi[J_n(2)Y_m(2) - J_m(2)Y_n(2)]. \quad (12)$$

This function inherits recursion (11) in parameter n with $x = 2$, because for m fixed Υ is a linear combination of $J_n(2)$ and $Y_n(2)$, so that

$$\Upsilon(n+1, m) + \Upsilon(n-1, m) = n\Upsilon(n, m). \quad (13)$$

Claim 3. *For all integers n and m the function $\Upsilon(n, m)$ is integer valued.*

Proof. Clearly $\Upsilon(m, m) = 0$ so it suffices to show that $\Upsilon(m+1, m)$ is integer valued, since recursion (13) takes care of all other values.

Two important relations for the Bessel functions are needed. The first is that

$$xJ'_n(x) = nJ_n(x) - xJ_{n+1}(x), \quad \forall n. \quad (14)$$

and the same holds with Y_n replacing J_n , see 3.2(4), p. 45 and 3.56(4), p. 66 of [Wat22], resp. The second is a fact relating to the Wronskian, namely that

$$\frac{2}{\pi x} = J_n(x)Y'_n(x) - J'_n(x)Y_n(x), \quad \forall n, x \neq 0, \quad (15)$$

see 3.63(1), p. 76 of [Wat22]. By dividing both sides of equation (14) by $x = 2$ and using the resulting expression for $J'_n(2)$ and $Y'_n(2)$, resp., in (15) results in

$$\begin{aligned} \frac{1}{\pi} &= J_n(2) \left[\frac{n}{2}Y_n(2) - Y_{n+1}(2) \right] - \left[\frac{n}{2}J_n(2) - J_{n+1}(2) \right] Y_n(2) \\ &= J_{n+1}(2)Y_n(2) - J_n(2)Y_{n+1}(2) = \frac{1}{\pi}\Upsilon(n+1, n), \end{aligned}$$

the last equality by the definition (12) of Υ . Hence, $\Upsilon(n+1, n) = 1$. \square

Now we can evaluate the function $\Upsilon(n, m)$ for different values of n and m to find that $B_n = \Upsilon(n+2, 0)$, $n = 2, 3$, and $A_n = 2\Upsilon(n+2, 3) + \Upsilon(n+2, 0)$, $n = 2, 3$, see Table 2. Since $\Upsilon(n+2, m)$ satisfies recursion (10) in n this is enough to know that they agree for all $n \geq 2$.

By relation (8), definition (12) and the above expression for A_n and B_n we get

$$\begin{aligned} b_n &= \Upsilon(n+3, 0) - \Upsilon(n+2, 0) \quad \text{and} \\ a_n &= 2[\Upsilon(n+3, 3) - \Upsilon(n+2, 3)] + \Upsilon(n+3, 0) - \Upsilon(n+2, 0). \end{aligned} \quad (16)$$

n	1	2	3	4	5	6	7
b_n	1	5	26	158	1113	8937	80624
$B_n = (b_n - b_{n-1})/n$		2	7	33	191	1304	10241
$\Upsilon(n, 0)$	1	1	1	2	7	33	191
a_n	3	11	56	340	2395	19231	173490
$A_n = (a_n - a_{n-1})/n$		4	15	71	411	2806	22037
$2\Upsilon(n, 3) + \Upsilon(n, 0)$	-3	-1	1	4	15	71	411

Table 2: Connection between parameters a_n, b_n and $\Upsilon(n, m)$.

Henceforth we abbreviate $J_n = J_n(2)$ and $Y_n = Y_n(2)$, hence e.g.

$$J_n = \sum_{k=0}^{\infty} \frac{(-1)^k}{k!(k+n)!}.$$

We need asymptotics for a_n and b_n .

Claim 4.

$$\lim_{n \rightarrow \infty} \frac{b_n}{(n+2)!} = J_0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{a_n}{(n+2)!} = 2J_3 + J_0. \quad (17)$$

Proof. [AS64] lists the following asymptotic relations, for fixed x and n tending to infinity,

$$J_n(x) \sim \frac{1}{\sqrt{2\pi n}} \left(\frac{ex}{2n}\right)^n \quad \text{and} \quad Y_n(x) \sim -\sqrt{\frac{2}{\pi n}} \left(\frac{2n}{ex}\right)^n,$$

hence, by the well-known Stirling formula $n! \sim \sqrt{2\pi n} n^{n+1/2} e^{-n}$ we conclude that $J_n \sim 1/n!$ and $Y_n \sim -(n-1)!/\pi$. Thus

$$\Upsilon(n, m) \sim \pi \frac{1}{n!} Y_m + \pi J_m \frac{(n-1)!}{\pi} \sim J_m \cdot (n-1)!,$$

from which (17) follows. \square

Now we are ready to find an expression for π_0 . Since $\pi_n \rightarrow 0$ as $n \rightarrow \infty$ we get

$$\pi_0 = \lim_{n \rightarrow \infty} \frac{\pi_n + b_n}{a_n} = \frac{J_0}{2J_3 + J_0} = 0.4647184275\dots \quad (18)$$

This in turn gives us

$$\begin{aligned}
\pi_n &= a_n \pi_0 - b_n \\
&= 2[\Upsilon(n+3, 3) - \Upsilon(n+2, 3)]\pi_0 + [\Upsilon(n+3, 0) - \Upsilon(n+2, 0)][\pi_0 - 1] \\
&= \frac{2\pi}{2J_3 + J_0} \left([J_{n+3}Y_3 - J_3Y_{n+3} - J_{n+2}Y_3 + J_3Y_{n+2}]J_0 \right. \\
&\quad \left. - [J_{n+3}Y_0 - J_0Y_{n+3} - J_{n+2}Y_0 + J_0Y_{n+2}]J_3 \right) \\
&= \frac{2\pi(J_3Y_0 - J_0Y_3)}{2J_3 + J_0}(J_{n+2} - J_{n+3}) = \frac{2\Upsilon(3, 0)}{2J_3 + J_0}(J_{n+2} - J_{n+3}) \\
&= \frac{2}{2J_3 + J_0}(J_{n+2} - J_{n+3}), \quad n \geq 1. \tag{19}
\end{aligned}$$

Then $\sum_0^N \pi_j = 1 - 2J_{N+3}/(2J_3 + J_0)$ and as $J_n \rightarrow 0$, as $n \rightarrow \infty$, we can verify that Π is indeed a distribution.

2.2 The time constant

Following the discussion before and after (1) we get the time constant

$$\tau = \frac{1}{1 + \pi_0} = \frac{2J_3 + J_0}{2J_3 + 2J_0} = 0.6827250759 \dots \tag{20}$$

This can be compared with first-passage percolation on \mathbb{N} , i.e. the infinite path graph (with exponential r.v.'s with mean 1), where the time constant necessarily is 1. We may also compare it with $1/2$ which is the time constant if the rungs of the ladder were associated with random variables all identically 0, i.e. there is immediate infection between nodes on different levels at distance 1 from each other.

Note that we started the process by infecting both $(0, 0)$ and $(0, 1)$. Consider the infection of the single node $(0, 0)$ at time 0. Then, the front process \hat{F} will not be a continuous time Markov chain (as before), as the behaviour of the front is very different before and after the random time Z until there are infected nodes on both levels. Since we are interested in stationarity we may simply disregard what happens before Z and view this as starting the front process F at time Z , $F_t = \hat{F}_{Z+t}$, $t \geq 0$. Then F is a continuous time Markov chain with intensity matrix given by (2), as before, albeit started in a random state instead of $F_0 = 0$. Z is dominated by a mean 1 exponential random variable (the edge between $(0, 0)$ and $(0, 1)$). As F will converge to the stationary distribution Π , so will \hat{F} .

Hence, infecting just a single node initially makes no difference to the time constant. (In fact, starting with an initial infection of any finite number of nodes will not change the time constant.)

This time constant also gives an upper bound for the time constant for first-passage percolation with mean 1 exponential r.v.'s on \mathbb{Z}^2 , but not stronger than already existing bounds, see [AP02].

2.3 Residual times

The fact that we can write down the stationary distribution Π for the process F_t also gives us the opportunity to calculate another statistic for first passage percolation on the ladder, namely the average residual time. As far as we know, this quantity has not previously been studied for percolation. Define the *residual time at t* as $R_t = \inf\{s : N_{t+s} = N_t + 1\}$, i.e. the time it takes, after t , for the infection to spread one more step up the ladder. We are interested in $T = \mathbb{E}R_t$ for a late time t .

One might a priori think that T and τ are the same thing. If we were looking at percolation on \mathbb{N} , which is the same as a Poisson process, then they are. This is the waiting time paradox; on one hand: looking at a fixed time point increases the probability of choosing a long time interval, but on the other: typically half that interval has passed. For a Poisson process, these effects balance out. For percolation on a ladder, as shown in (20) and (23) below, they do not.

Now, to calculate T we will assume that F has its stationary distribution, so that we may disregard time and put $R = R_t$. Let $\gamma_n = \mathbb{E}[R|F = n]$. Hence, γ_n is the expected time it takes for the infection to spread another step given that the front is in state n .

If $F = 0$ then the residual time is the minimum of two mean 1 exponential random variables, so that $\gamma_0 = 1/2$. For $n \geq 1$, we write down a recursive formula for γ_n . If $F = n$ then there are $n + 2$ possible edges, equally likely, that the infection might spread along. Only one of these results in an increase in the height of the infection, namely the one that gives $F = n + 1$. Two edges result in $F = n - 1$ and the remaining ones to one of the states $\{0, 1, \dots, n - 2\}$, respectively. This gives the formula

$$\gamma_n = \frac{1}{n+2} \left(1 + 2\gamma_{n-1} + \sum_{j=0}^{n-2} \gamma_j \right). \quad (21)$$

By writing down the formula for γ_{n-1} one sees that

$$1 + \sum_0^{n-2} \gamma_j = (n+1)\gamma_{n-1} - \gamma_{n-2}$$

which inserted into (21) yields the incremental relation

$$(n+2)(\gamma_n - \gamma_{n-1}) = \gamma_{n-1} - \gamma_{n-2},$$

which together with the fact that $\gamma_1 = \frac{1}{3}(1 + 2\frac{1}{2}) = \frac{2}{3}$ in turn gives us the formulas

$$\gamma_n - \gamma_{n-1} = \frac{1}{(n+2)!}, \quad n \geq 1, \quad \text{and} \quad \gamma_n = \sum_{j=0}^{n+2} \frac{1}{j!}, \quad n \geq 0. \quad (22)$$

Then, as F_t is in state j with probability π_j where it remains on average γ_j amount of time before N_t increases, we get the average residual time T as

$$\begin{aligned} T &= \sum_{n=0}^{\infty} \pi_n \gamma_n = \frac{1}{2J_3 + J_0} \left(J_0/2 + 2 \sum_{n=1}^{\infty} (J_{n+2} - J_{n+3}) \gamma_n \right) \\ &= \frac{1}{2J_3 + J_0} \left(\frac{1}{2} J_0 + 2J_3 \gamma_1 + 2 \sum_{n=1}^{\infty} (\gamma_{n+1} - \gamma_n) J_{n+3} \right) \\ &= \frac{1}{2J_3 + J_0} \left(\frac{1}{2} J_0 + \frac{4}{3} J_3 + 2 \sum_{n=1}^{\infty} \frac{J_{n+3}}{(n+3)!} \right) = 0.5953444665 \dots \quad (23) \end{aligned}$$

Acknowledgements: Thanks to Sven Erick Alm for introducing me to this problem and to Svante Janson for making useful suggestions regarding the mathematics. Also, we wish to thank an anonymous referee for carefully reading the manuscript and providing us with numerous helpful comments.

References

- [AS64] M. Abramowitz, I. Stegun: *Handbook of mathematical functions*. National Bureau of Standards Applied Mathematics Series 55 (1964).
- [AJ] S. E. Alm, S. Janson; personal communication.
- [AJ90] S. E. Alm, S. Janson: *Random self-avoiding walks on one-dimensional lattices*. Commun. Statist. -Stochastic Models, **6**, No. 2 (1990), 169–212.
- [AP02] S.E. Alm, R. Parviainen: *Lower and Upper Bounds for the Time Constant of First-Passage Percolation*. Combinatorics, Probability and Computing **11**, No. 5 (2002) 433–445
- [FGS06] A. Flaxman, D. Gamarnik, G. Sorkin: *First-passage percolation on a width-2 strip and the path cost in a VCG auction*. Internet and Network Economics **4286** (2006) 99–111.
- [PR] R. Parvianen, H. Renlund. In preparation.
- [Sch09] E. Schlemm: *First-passage percolation rates on width-two stretches with exponential link weights*, Electron. Commun. Probab., **14** (2009), 424–434.
- [Sl07] N. Sloane: *The On-Line Encyclopedia of Integer Sequences*. Published electronically at www.research.att.com/~njas/sequences/ (2007).

[SW78] R. T. Smythe, J. C. Wierman: *First-Passage Percolation on the square lattice*. Lecture notes in mathematics **671**, Springer (1978).

[Wat22] G. N. Watson: *A Treatise on the Theory of Bessel Functions*. Cambridge University Press (1922).