

## Homework I- Liese Course II deadline: 3/11/09

1. Consider two models

$$\mathcal{M} = (\mathcal{X}, \mathfrak{A}, \{P_\theta\}_{\theta \in \Theta}); \mathcal{N} = (\mathcal{Y}, \mathfrak{B}, \{P_\theta^{T(\mathbf{X})}\}_{\theta \in \Theta})$$

Suppose  $T(\mathbf{X})$  is a sufficient statistic for  $\theta \in \Theta$ . Show that the models are equivalent.

2. Consider two models  $\mathcal{M}, \mathcal{N}$  and two equivalent models of them  $\mathcal{M}_{suff}, \mathcal{N}_{suff}$  with  $\Delta(\mathcal{M}, \mathcal{M}_{suff}) = 0, \Delta(\mathcal{N}, \mathcal{N}_{suff}) = 0$ . Show

$$\Delta(\mathcal{M}, \mathcal{N}) = \Delta(\mathcal{M}_{suff}, \mathcal{N}_{suff})$$

3. Consider two models

$$\mathcal{M}_1 = (\mathbb{R}, \mathfrak{B}, \{N(\theta, \sigma_1^2)\}_{\theta \in \mathbb{R}}); \mathcal{M}_2 = (\mathbb{R}, \mathfrak{B}, \{N(\theta, \sigma_2^2)\}_{\theta \in \mathbb{R}})$$

Derive an upper bound for  $\Delta(\mathcal{M}_1, \mathcal{M}_2)$ .

4. Consider two models

$$\mathcal{M}_1 = (\{0, 1\}^N, \mathfrak{B}, \{Ber(p)^{\otimes N}\}_{p \in (0,1)}); \mathcal{M}_2 = (\{0, 1\}^N, \mathfrak{B}, \{Ber(\pi p)^{\otimes N}\}_{p \in (0,1)})$$

for some  $\pi \in (0, 1)$ . Derive an upper bound for  $\Delta(\mathcal{M}_1, \mathcal{M}_2)$ .

5. Compare the models

$$\begin{aligned} \mathcal{M}_1 &= (\{0, 1, \dots, n\}, \mathcal{P}(\{0, 1, \dots, n\}), \{Bin(n, p)\}_{p \in (0,1)}) \\ \mathcal{M}_2 &= (\{0, 1, \dots, n\}^3, \mathcal{P}(\{0, 1, \dots, n\}^3), \{multinomial(n, \frac{p}{2}, \frac{p}{2}, (1-p))\}_{p \in (0,1)}). \end{aligned}$$

6. Compare the models

$$\begin{aligned} \mathcal{M}_1 &= (\{0, 1, \dots, n\}, \mathcal{P}(\{0, 1, \dots, n\}), \{Bin(n, p)\}_{p \in (0,1)}) \\ \mathcal{M}_2 &= (\{0, 1, \dots, n\}^3, \mathcal{P}(\{0, 1, \dots, n\}^3), \{multinomial(n, \frac{p}{3}, \frac{p}{3}, (1 - \frac{2p}{3}))\}_{p \in (0,1)}). \end{aligned}$$