

Analysis for PhD students (2025); Assignment 2

Problem 1. Prove that if $f \in L^1(\mathbb{R})$ and $\int_{-\infty}^{\infty} |\xi \widehat{f}(\xi)| d\xi < \infty$, then f coincides almost everywhere with a differentiable function f_0 whose derivative is

$$f_0'(x) = 2\pi i \int_{-\infty}^{\infty} \xi \widehat{f}(\xi) e^{2\pi i \xi x} d\xi \quad (\forall x \in \mathbb{R}). \quad (15p)$$

Problem 2. Prove that if μ is a positive Borel measure on \mathbb{T} with $\mu(\mathbb{T}) = 1$, then $|\widehat{\mu}(k)| < 1$ for all $k \neq 0$ unless μ is a linear combination, with nonnegative coefficients, of the point masses at $y, y + \frac{1}{m}, \dots, y + \frac{m-1}{m}$ for some $m \in \mathbb{N}$ and $y \in \mathbb{T}$. (15p)

Problem 3. Folland p. 255, Exercise 15, all parts a–c (the Sampling Theorem). (15p)

Problem 4. Prove that for any $k \geq 1$:

$$\int_{-\infty}^{\infty} \frac{dx}{(1+|x-a|)^k(1+|x|)^k} \asymp_k \begin{cases} (1+|a|)^{-k} & \text{if } k > 1 \\ (1+|a|)^{-1} \log(2+|a|) & \text{if } k = 1, \end{cases}$$

uniformly over all $a \in \mathbb{R}$. (10p)

Problem 5. Let a_0 be a fixed number with $0 < a_0 < 1$. Prove that

$$\int_1^{\infty} e^{-bx} x^{ab} dx \sim \frac{e^{-b}}{(1-a)b}$$

as $b \rightarrow \infty$, uniformly over all $a \in [0, a_0]$. (15p)

Problem 6. Let $n \in \mathbb{Z}^+$. Find (with proof) the largest possible constant $\alpha > 0$ such that

$$\int_0^1 e(x_1 t + x_2 t^2 + \cdots + x_n t^n) dt \ll_n (1 + |x|)^{-\alpha}, \quad \forall x \in \mathbb{R}^n. \quad (15p)$$

Problem 7. Using stationary phase, prove that for any fixed $\alpha > 1$,

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} e^{i\lambda(\alpha \sin x - x)} dx = \sqrt{\frac{2}{\pi}} \cdot \frac{\cos\left(\lambda(\sqrt{\alpha^2 - 1} - \arccos(\alpha^{-1})) - \frac{\pi}{4}\right)}{\sqrt[4]{\alpha^2 - 1} \sqrt{\lambda}} + O\left(\frac{1}{\lambda}\right)$$

as $\lambda \rightarrow +\infty$ (through real numbers), where the implied constant may depend on α .

[Comment: Specializing to $\lambda = n \in \mathbb{Z}^+$ this gives an asymptotic formula for $J_n(\alpha n)$, via (8.9) in the lecture notes. It is satisfactory to note that this asymptotic formula is consistent with Prop. 10.1 in the lecture notes.]

(15p)

To be returned: Wednesday, November 19, before midnight. Please send your solutions by email, or put them in my mailbox.