

RADEMACHER CHAOS: TAIL ESTIMATES VS LIMIT THEOREMS

RON BLEI AND SVANTE JANSON

ABSTRACT. We study Rademacher chaos indexed by a sparse set which has a fractional combinatorial dimension. We obtain tail estimates for finite sums and a normal limit theorem as the size tends to infinity. The tails for finite sums may be much larger than the tails of the limit.

1. INTRODUCTION AND RESULTS

A (homogeneous) Rademacher chaos is a random variable of the type

$$S = \sum_{i_1 < \dots < i_d} a_{i_1 \dots i_d} r_{i_1} \cdots r_{i_d}, \quad (1.1)$$

where $d \geq 1$, $a_{i_1 \dots i_d}$ are real or complex numbers and r_1, r_2, \dots is a sequence of independent random variables with the symmetric two-point distribution $\mathbb{P}(r_i = 1) = \mathbb{P}(r_i = -1) = 1/2$. (For example, r_i could be the classical Rademacher functions [13], defined on $[0, 1]$ (with the usual Lebesgue measure) by $r_i(x) = 1 - 2b_i$ when $x \in [0, 1]$ has the binary expansion $0.b_1 b_2 \dots$, but it is often more convenient to let r_i be defined on the Cantor group \mathbb{Z}_2^∞ . For our purposes, the choice of r_i does not matter.) Equivalently, S is a linear combination of the Walsh functions of the type $r_{i_1} \cdots r_{i_d}$.

We will consider only finite sums (1.1), so there is no problem of convergence, and all moments of S are finite.

We are interested in two related properties of the random variables S : the tail behaviour, i.e. the size of the probabilities $\mathbb{P}(|S| > x)$ for large x , and the size of the L^q norms $\|S\|_q = (\mathbb{E} |S|^q)^{1/q}$ for large q . For convenience, we define $\tilde{S} = S/\|S\|_2$; thus $\mathbb{E} \tilde{S} = 0$ and $\text{Var} \tilde{S} = \mathbb{E} \tilde{S}^2 = 1$.

Bonami's hypercontractive inequality [3] implies that every S in (1.1) satisfies

$$\|S\|_q \leq (q-1)^{d/2} \|S\|_2 = (q-1)^{d/2} \left(\sum_{i_1 < \dots < i_d} |a_{i_1 \dots i_d}|^2 \right)^{1/2}, \quad q \geq 2, \quad (1.2)$$

or, equivalently, $\|\tilde{S}\|_q \leq (q-1)^{d/2}$, $q \geq 2$. (See also [1, 2, 7, 9].)

In general, this estimate is best possible, up to a constant depending on d but not on q . For example, it is easily seen that $a_{i_1 \dots i_d} = 1$ for $1 \leq i_1 < \dots < i_d \leq n$ (and 0 otherwise) yields an $S = S_n$ that after suitable normalization converges, as $n \rightarrow \infty$, to a (Hermite) d -degree polynomial in a Gaussian random variable; see Example 3.2. It thus follows that for some $c(d) > 0$ and every $q \geq 2$,

Date: May 29, 2002; revised September 24, 2002.

$\|S\|_q \geq c(d)(q-1)^{d/2}\|S\|_2$ provided n is large enough. (See e.g. [7, Chapter XIII].)

In this paper we study Rademacher chaos (1.1) where most coefficients $a_{i_1 \dots i_d} = 0$ so that we really only sum over an indexing set which is combinatorially sparse in the sense of [2, Chapters XII and XIII]. In this case, Bonami's hypercontractive inequality (1.2) can be improved, precisely reflecting the sparsity of the indexing set.

We first recall some definitions [2, Chapters XIII], which we modify and adapt to our purposes in this paper.

For $F \subseteq \mathbb{N}^d$ and $\alpha > 0$, define

$$\Psi_F(s) = \max\{|F \cap (A_1 \times \dots \times A_d)| : A_j \subseteq \mathbb{N}, |A_j| \leq s, j = 1, \dots, d\},$$

and

$$d_F(\alpha) = \sup_{s \geq 1} \Psi_F(s)/s^\alpha = \sup_{A_1, \dots, A_d} |F \cap (A_1 \times \dots \times A_d)| / (\max_{j \leq d} |A_j|)^\alpha.$$

In [2, §XIII.4], the combinatorial dimension of a set $F \subseteq \mathbb{N}^d$ is defined to be

$$\begin{aligned} \dim(F) &= \limsup_{s \rightarrow \infty} (\log \Psi_F(s) / \log s) \\ &= \sup\{\alpha : d_F(\alpha) = \infty\} = \inf\{\alpha : d_F(\alpha) < \infty\}. \end{aligned} \quad (1.3)$$

In this paper, we consider sequences of index sets $F_N \subseteq [N]^d$, where $[N] = \{1, \dots, N\}$, and we adopt the definition below. Since we only want to consider non-empty index sets, we consider sequences starting at some index $N_0 \geq 1$; this allows for some empty F_N for smaller N that we ignore.

Definition. A sequence $F_N \subseteq [N]^d$, $N = N_0, N_0 + 1, \dots$, has combinatorial dimension α if there exist positive constants C_1, C_2 such that for all $N \geq N_0$,

$$d_{F_N}(\alpha) \leq C_1,$$

(i.e. $|F_N \cap (A_1 \times \dots \times A_d)| \leq C_1 (\max_{j \leq d} |A_j|)^\alpha$), and

$$|F_N| \geq C_2 N^\alpha.$$

We write $\dim\{F_N\} = \alpha$.

Given a set $F \subseteq \mathbb{N}^d$, we define $F_N = F \cap [N]^d$. In the present paper, we define $\dim(F) = \dim\{F_N\}$ when the latter exists (and leave the dimension undefined otherwise).

Remark 1.1. Note that this is a stricter definition than (1.3); there are sets F with no dimension in the present sense, but it is easily seen that when the dimension exists in the present sense, it coincides with (1.3).

If the cardinalities of F_N are uniformly bounded, then $\dim\{F_N\} = 0$; otherwise $1 \leq \dim\{F_N\} \leq d$ (if $\dim\{F_N\}$ exists at all). We are mainly interested in the case $1 < \dim\{F_N\} < d$.

Let $\Delta^d = \{(i_1, \dots, i_d) : 1 \leq i_1 < \dots < i_d < \infty\}$ and $\Delta_N^d = \Delta^d \cap [N]^d$. We will in the sequel consider only $F \subseteq \Delta^d$ and $F_N \subseteq \Delta_N^d$; This is not essential, but restriction to ordered sets of indices is convenient when we study sums (1.6).

It is proved in [2, Chapter XIII] that for every $\alpha \in [1, d]$, there exist sets $F \subset \Delta^d$ of combinatorial dimension α (also in the stricter sense used here). Such sets can always be constructed by a random procedure; for rational $\alpha \geq 1$ and d such that $d\alpha$ is an integer, it is also possible to use the following deterministic construction.

Example 1.2. (*Minimal fractional Cartesian products* [2, §XIII.1 and p. 493].) Fix arbitrary integers $d \geq 3$ and $1 \leq m \leq d$, and let $\{S_1, \dots, S_d\}$ be a cover of $[d]$ consisting of m -subsets of $[d]$, such that every $i \in [d]$ appears in exactly m elements of S_1, \dots, S_d ; i.e., $\bigcup_{j=1}^d S_j = [d]$, $|S_j| = m$, and for every $i \in [d]$, $|\{j : i \in S_j\}| = m$.

We employ the following notation: if X is a set, $\mathbf{y} = (y_1, \dots, y_d) \in X^d$ and $S \subseteq [d]$, then

$$\pi_S \mathbf{y} = (y_i : i \in S).$$

For an integer $N \geq d^m$, let n be the greatest integer such that $n \leq N^{1/m}$. Fix a one-one map φ from $[n]^m$ into $[N]$, and consider

$$F_N^* = \{(\varphi(\pi_{S_1} \mathbf{k}), \dots, \varphi(\pi_{S_d} \mathbf{k})) : \mathbf{k} \in [n]^d\}. \quad (1.4)$$

In order to obtain a subset of Δ_N^d , for the purposes of this paper, we modify this set to

$$F_N = \{(i_1, \dots, i_d) \in \Delta_N^d : (i_{\rho_1}, \dots, i_{\rho_d}) \in F_N^* \text{ for some permutation } \rho\}. \quad (1.5)$$

We call the sequence $\{F_N\}$ a *fractional Cartesian product*.

We further say the fractional Cartesian product is *disconnected* if $[d]$ can be partitioned into two disjoint nonempty subsets T_1 and T_2 such that each S_j is a subset of either T_1 or T_2 , and *connected* otherwise.

The archetypal case is $d = 3$, $m = 2$, $S_1 = \{1, 2\}$, $S_2 = \{1, 3\}$ and $S_3 = \{2, 3\}$. This gives a connected fractional Cartesian product.

Claim. $\dim\{F_N\} = \dim\{F_N^*\} = d/m$.

We verify the claim in the archetypal case $d = 3$, $m = 2$ only. The general case is similar; see [2, Corollary XIII.16].

Let $1 \leq s \leq N$ be an integer, and let A, B, C be arbitrary subsets of $[N]$. Then,

$$|F_N^* \cap (A \times B \times C)| = \sum_{k_1, k_2, k_3 \in [n]} \mathbf{1}_A(\varphi(k_1, k_2)) \mathbf{1}_B(\varphi(k_1, k_3)) \mathbf{1}_C(\varphi(k_2, k_3)).$$

A three-fold application of the Cauchy–Schwarz inequality yields

$$\begin{aligned} |F_N^* \cap (A \times B \times C)| &\leq \left(\sum_{k_1, k_2 \in [n]} \mathbf{1}_A(\varphi(k_1, k_2)) \right)^{1/2} \left(\sum_{k_1, k_3 \in [n]} \mathbf{1}_B(\varphi(k_1, k_3)) \right)^{1/2} \\ &\quad \cdot \left(\sum_{k_2, k_3 \in [n]} \mathbf{1}_C(\varphi(k_2, k_3)) \right)^{1/2} \\ &\leq |A|^{1/2} |B|^{1/2} |C|^{1/2}, \end{aligned}$$

which implies $\Psi_{F_N^*}(s) \leq s^{3/2}$ and thus $\Psi_{F_N}(s) \leq 6s^{3/2}$. In the opposite direction,

$$|F_N| = |\Delta_n^3| = \binom{n}{3} \geq c_1 n^3 \geq c_2 N^{3/2}.$$

Remark 1.3. Again, the definition differs slightly from [2]; there the fractional Cartesian product is defined on an infinite set ($n = \infty$ in (1.4)).

Remark 1.4. Note that the function φ appears in the definition of a fractional Cartesian product, only because we let the indices be integers in this paper. We might avoid φ by changing the notation slightly; for example, for the case $d = 3$, $m = 2$, we could equivalently write (1.9) below as $S_N = \sum_{i < j < k \leq n} r_{ij} r_{ik} r_{jk}$, where r_{ij} , $i < j$ are independent Rademacher variables.

It is shown in [2] (e.g., Corollary XIII.8.29) that if $F \subseteq \Delta^d$ (finite or infinite), and S is a Rademacher chaos

$$S = \sum_{(i_1, \dots, i_d) \in F} a_{i_1 \dots i_d} r_{i_1} \cdots r_{i_d}, \quad (1.6)$$

then

$$\|S\|_q \leq K d_F(\alpha)^{1/2} q^{\alpha/2} \|S\|_2, \quad q \geq 1, \quad (1.7)$$

where $K < \infty$ depends only on the ambient dimension d . (See Section 5 below.) In particular, if $\dim\{F_N\} < d$, the exponent in (1.2) can be improved, with d replaced by the combinatorial dimension.

These norm estimates lead to tail estimates by the customary procedure: If (1.7) holds and $d_F(\alpha) < \infty$, then for any $x > 0$ and $q \geq 1$, by Markov's inequality,

$$\mathbb{P}(|\tilde{S}| \geq x) \leq x^{-q} \mathbb{E} |\tilde{S}|^q = x^{-q} \|\tilde{S}\|_q^q \leq (x^{-1} C q^{\alpha/2})^q,$$

where $C = K d_F(\alpha)^{1/2}$. Taking $q = (x/C)^{2/\alpha} e^{-1}$ (if $x \geq C e^{\alpha/2}$), we obtain

$$\mathbb{P}(|\tilde{S}| \geq x) \leq e^{-\alpha q/2} = \exp(-c x^{2/\alpha}), \quad (1.8)$$

for a constant $c > 0$ depending on d , α and $d_F(\alpha)$ only.

The norm and tail estimates above are in fact sharp, in a sense made precise below. (Cf. [2, Corollary XIII.8.29].) For simplicity, we will consider only the case where $a_{i_1 \dots i_d} = 1$ or 0 . Specifically, we consider a sequence of non-empty sets $F_N \subseteq \Delta_N^d$ and Rademacher chaos

$$S_N = \sum_{(i_1, \dots, i_d) \in F_N} r_{i_1} \cdots r_{i_d}. \quad (1.9)$$

Clearly, $\|S_N\|_2 = |F_N|^{1/2}$, and thus $\tilde{S}_N = |F_N|^{-1/2} S_N$.

Theorem 1.5. *Suppose $\dim\{F_N\} = \alpha \geq 1$, where $F_N \subseteq \Delta_N^d$. Let S_N be given by (1.9). Then there exist positive constants c_1, c_2, c_3, c_4 such that for every $q \geq 1$,*

$$c_1 q^{\alpha/2} \leq \sup_N \|\tilde{S}_N\|_q \leq c_2 q^{\alpha/2}, \quad (1.10)$$

and for all $x \geq 2$,

$$\exp(-c_3 x^{2/\alpha}) \leq \sup_N \mathbb{P}(|\tilde{S}_N| > x) \leq \exp(-c_4 x^{2/\alpha}). \quad (1.11)$$

A natural question arises: can \sup_N in (1.10) and (1.11) be replaced by $\lim_{N \rightarrow \infty}$? (See Remark ii in [2, p.524].) In the standard integer-dimensional case $F_N = \Delta^d$, the answer is affirmative (by a d -fold application of the usual Central Limit theorem). But in many fractional-dimensional cases, the answer is negative: the precise relation between tail estimates and combinatorial dimension, as per (1.11), is completely wiped out in the limit. We illustrate this in two important cases.

Theorem 1.6. *Let $F_N \subseteq \Delta_N^d$, $N = 1, \dots$, and let S_N be given by (1.9). Suppose either (i) $d = 2$ and $1 < \dim\{F_N\} < 2$, or (ii) F_N is a connected fractional Cartesian product as in Example 1.2. Then $\tilde{S}_N \xrightarrow{d} N(0, 1)$ with convergence of all moments. In particular, if $\xi \sim N(0, 1)$, then, for all $q \geq 1$,*

$$\lim_{N \rightarrow \infty} \|\tilde{S}_N\|_q = \|\xi\|_q \leq q^{1/2}$$

and for all $x \geq 2$,

$$\lim_{N \rightarrow \infty} \mathbb{P}(|\tilde{S}_N| > x) = \mathbb{P}(|\xi| > x) \leq \exp(-x^2/2).$$

Case (ii) with $m = 2$ can be translated (using Remark 1.4) into a result for random graphs, which is a special case (with $p = 1/2$) of [5, Theorem 1]; see also [6] and [7, Chapter XI].

Theorems 1.5 and 1.6 complement one another in the following (heuristic) sense. Let us agree that tail probabilities of sums of uncorrelated symmetric variables provide a gauge of interdependence between the variables: larger tail probabilities (smaller likelihood of cancellations) convey higher degree of interdependence, and conversely. In this light, Theorem 1.5 provides a precise assessment of interdependence of the random variables $r_{i_1} \cdots r_{i_d}$, $(i_1, \dots, i_d) \in F_N$. As a counterpoint, reflecting increasing sparsity of F_N relative to the full product set $\Delta_N^{(d)}$, Theorem 1.6 asserts that F_N in the limit, as $N \rightarrow \infty$, is asymptotically independent.

Theorems 1.5 and 1.6 show that, for large q or x , the limits as $N \rightarrow \infty$ are much smaller than the largest values for finite N . If we fix a large q and study $\|\tilde{S}_N\|_q$ as N grows, we begin with rather small values (at most $|F_N|^{1/2}$) that grow to a maximum of the order $q^{\alpha/2}$ (when N is about q , see Section 2), but then the norms decrease again towards a limit of the order $q^{1/2}$. (We do not know whether the increase and decrease are monotone; there might be several local maxima.) A similar story holds for $\mathbb{P}(|\tilde{S}_N| > x)$ for a fixed large x . Consequently, the limit results in Theorem 1.6 are misleading when we consider \tilde{S}_N for finite N .

A central limit theorem in fact holds generally under a condition of sparsity in F_N that is milder than the sparsity implied by non-integer combinatorial dimension. The condition is in effect that F_N is not “too close” to a product

set. To express this precisely we use the following terminology. For $j \in [N]$, define

$$F_{Nj}^* = \{(i_1, \dots, i_d) \in F_N : j \in \{i_1, \dots, i_d\}\}.$$

Further, let $F_N^\#$ be the subset of $F_N \times F_N$ defined as follows: a pair of d -tuples $((i_1, \dots, i_d), (j_1, \dots, j_d)) \in F_N^\#$ if $\{i_1, \dots, i_d\} \cap \{j_1, \dots, j_d\} = \emptyset$ and there exist $(k_1, \dots, k_d) \in F_N$, $(l_1, \dots, l_d) \in F_N$ such that $\{k_1, \dots, k_d, l_1, \dots, l_d\} = \{i_1, \dots, i_d, j_1, \dots, j_d\}$ but (k_1, \dots, k_d) does not equal (i_1, \dots, i_d) or (j_1, \dots, j_d) . (In other words, the $2d$ indices $i_1, \dots, i_d, j_1, \dots, j_d$ can be partitioned in at least two ways into elements of F_N .)

Theorem 1.7. *Suppose*

$$\lim_{N \rightarrow \infty} \max_j |F_{Nj}^*|/|F_N| = 0 \quad (1.12)$$

and

$$\lim_{N \rightarrow \infty} (|F_N^\#|/|F_N|^2) = 0. \quad (1.13)$$

Then $\tilde{S}_N \xrightarrow{d} N(0, 1)$, with convergence of all moments.

We have the following partial converse. (The trivial example $F = \{(1, j) : j \geq 2\}$ shows that (1.12) is not necessary; we do not know whether it is needed at all in Theorem 1.7.)

Theorem 1.8. *Suppose that $\tilde{S}_N \xrightarrow{d} N(\mu, \sigma^2)$ for some μ and $\sigma^2 > 0$. Then $\mu = 0$, $\sigma^2 = 1$ and (1.13) holds.*

The proof of Theorem 1.5 is given in Section 2, and the proofs of Theorems 1.6, 1.7, 1.8 are given in Section 3. Some simple examples of nonnormal limits when (1.13) is not satisfied are given also in Section 3. Further remarks and open problems are presented in Section 4. Finally, Section 5 contains corrections to arguments in [2] that are relevant to this paper.

2. PROOF OF THEOREM 1.5

The upper bounds follow by (1.7) and (1.8) (for $x \geq x_0$, say; the case $2 \leq x \leq x_0$ follows by Chebyshev's inequality if c_4 is small enough).

To verify the lower bounds, let \mathcal{E}_N be the event $r_1 = \dots = r_N = 1$; thus $\mathbb{P}(\mathcal{E}_N) = 2^{-N}$. On \mathcal{E}_N , we have $S_N = |F_N|$ and thus $\tilde{S}_N = |F_N|^{1/2}$. Hence, for every $q \geq 1$,

$$\|\tilde{S}_N\|_q \geq |F_N|^{1/2} \mathbb{P}(\mathcal{E}_N)^{1/q} \geq cN^{\alpha/2} 2^{-N/q}.$$

To verify the left inequality in (1.10), in the line above choose $N = \max(N_0, \lfloor q \rfloor)$.

Similarly, given x (large enough), let $N = \lceil Cx^{2/\alpha} \rceil$ for a constant $C > C_2^{1/\alpha}$. Then, on \mathcal{E}

$$\tilde{S}_N = |F_N|^{1/2} \geq C_2^{1/2} N^{\alpha/2} > x,$$

and thus

$$\mathbb{P}(\tilde{S}_N > x) \geq \mathbb{P}(\mathcal{E}_N) = 2^{-N} \geq \exp(-c_3 x^{2/\alpha}). \quad \square$$

3. ASYMPTOTIC NORMALITY

Lemma 3.1. *If $d = 2$, then for any α and any finite subsets A and B of \mathbb{N} such that $|A| \leq |B|$,*

$$\begin{aligned} |F \cap (A \times B)| &\leq 2d_F(\alpha)|A|^{\alpha-1}|B|, \\ |F \cap (B \times A)| &\leq 2d_F(\alpha)|A|^{\alpha-1}|B|. \end{aligned}$$

Proof. We may assume $|A| \geq 1$. Partition B into $\lceil |B|/|A| \rceil \leq 2|B|/|A|$ subsets B_j with $|B_j| \leq |A|$. For each j , $|F \cap (A \times B_j)| \leq \Psi_F(|A|) \leq d_F(\alpha)|A|^\alpha$, and similarly for $|F \cap (B_j \times A)|$. The result follows by summing over j . \square

Proof of Theorem 1.6. We verify the conditions of Theorem 1.7. Let $\alpha = \dim\{F_N\}$.

First consider case (i), i.e., suppose that $d = 2$ and $1 < \alpha < 2$. Then, $|F_{Nj}^*| \leq N = o(|F_N|)$, which verifies (1.12).

Next, choose $\varepsilon_N > 0$ such that $\varepsilon_N \rightarrow 0$ and $N\varepsilon_N^{1/(2-\alpha)} \rightarrow \infty$. (For example, $\varepsilon_N = N^{-\delta}$ for $0 < \delta < 2 - \alpha$.) Let $A = A_N = \{i : |F_{Ni}^*| \geq \varepsilon_N N\}$. Then,

$$|F_N \cap (A \times [N])| + |F_N \cap ([N] \times A)| = \sum_{i \in A} |F_{Ni}^*| \geq \varepsilon_N N |A|,$$

and thus, by Lemma 3.1

$$\varepsilon_N N |A| \leq 4d_{F_N}(\alpha)|A|^{\alpha-1}N,$$

which implies

$$|A| \leq (4d_{F_N}(\alpha)/\varepsilon_N)^{1/(2-\alpha)} = o(N). \quad (3.1)$$

By definition, $F_N^\#$ is the set of all $((i, j), (k, l)) \in F_N \times F_N$, all of which entries are distinct, such that either $((i, k), (j, l)) \in F_N \times F_N$ or $((i, l), (j, k)) \in F_N \times F_N$ (or both). We let $F_{N1}^\#$ be the subset of $F_N^\#$ where $i \in A$, and $F_{N2}^\#$ the subset where $i \notin A$.

The number of possible $(i, j) \in F_N$ with $i \in A$ is $|F_N \cap (A \times [N])|$, and thus, by Lemma 3.1 and (3.1),

$$\begin{aligned} |F_{N1}^\#| &\leq |F_N \cap (A \times [N])| \cdot |F_N| \leq 2d_{F_N}(\alpha)|A|^{\alpha-1}N|F_N| \\ &= o(N^\alpha|F_N|) = o(|F_N|^2). \end{aligned} \quad (3.2)$$

On the other hand, let $F_{Ni}^{**} = \{k : (i, k) \in F_{Ni}^* \text{ or } (k, i) \in F_{Ni}^*\}$. Thus $|F_{Ni}^{**}| = |F_{Ni}^*|$. If $((i, j), (k, l)) \in F_N^\#$, then either k or l is in F_{Ni}^{**} , and thus the number of possible $(k, l) \in F_N$ for a given $i \notin A$ is at most, again by Lemma 3.1,

$$|F_N \cap (F_{Ni}^{**} \times [N])| + |F_N \cap ([N] \times F_{Ni}^{**})| \leq 4d_F(\alpha)|F_{Ni}^{**}|^{\alpha-1}N \leq 4d_F(\alpha)\varepsilon_N^{\alpha-1}N^\alpha.$$

Summing over all possible (i, j) we find

$$|F_{N2}^\#| \leq 4d_F(\alpha)\varepsilon_N^{\alpha-1}N^\alpha|F_N| = o(|F_N|^2). \quad (3.3)$$

Combining (3.2) and (3.3), we obtain (1.13) and the result follows in this case.

In case (ii), we first observe that fixing an index j in F_N means that the corresponding \mathbf{k} in (1.4) is such that $\pi_{S_i} \mathbf{k} = \varphi^{-1}(j)$ for some i ; for each i

this means that m of the d coordinates of \mathbf{k} have given values, so the number of choices of \mathbf{k} is at most dn^{d-m} . Consequently, $|F_{Nj}^*| \leq dn^{d-m} = o(|F_N|)$, proving (1.12).

Next, suppose that $((i_1, \dots, i_d), (j_1, \dots, j_d)) \in F_N^\#$, and that (i_1, \dots, i_d) and (j_1, \dots, j_d) are generated by (1.4) and (1.5) by some vectors \mathbf{i} and \mathbf{j} in $[n]^d$, respectively. By the definition of $F_N^\#$, there exists also $(k_1, \dots, k_d) \in F_N$, generated in the same way by, say, $\mathbf{k} \in [n]^d$, such that $\{k_1, \dots, k_d\} \subseteq \{i_1, \dots, i_d, j_1, \dots, j_d\}$ but (k_1, \dots, k_d) does not equal (i_1, \dots, i_d) or (j_1, \dots, j_d) .

Hence, each $\pi_{S_\nu} \mathbf{k}$, $1 \leq \nu \leq d$, coincides with some $\pi_{S_\mu} \mathbf{i}$ or $\pi_{S_\mu} \mathbf{j}$, $1 \leq \mu \leq d$. Define

$$\begin{aligned} J_1 &= \{\nu \in [d] : \pi_{S_\nu} \mathbf{k} = \pi_{S_\mu} \mathbf{i} \text{ for some } \mu\}; \\ J_2 &= \{\nu \in [d] : \pi_{S_\nu} \mathbf{k} = \pi_{S_\mu} \mathbf{j} \text{ for some } \mu\}; \\ T_s &= \bigcup_{\nu \in J_s} S_\nu, \quad s = 1, 2. \end{aligned}$$

Then $J_1 \cup J_2 = [d]$ and $T_1 \cap T_2 \neq \emptyset$, because otherwise the fractional Cartesian product would be disconnected.

If $q \in T_1$, then $q \in S_\nu$ for some $\nu \in J_1$, and thus $\pi_{S_\nu} \mathbf{k} = \pi_{S_\mu} \mathbf{i}$ for some μ . In particular, the q th coordinate of \mathbf{k} is one of the coordinates of \mathbf{i} . Similarly, if $q \in T_2$, then the q th coordinate of \mathbf{k} is one of the coordinates of \mathbf{j} .

Since $T_1 \cap T_2 \neq \emptyset$, this means that \mathbf{i} and \mathbf{j} have at least one coordinate in common (not necessarily in the same position). Consequently, the number of possible pairs (\mathbf{i}, \mathbf{j}) is $O(n^{2d-1})$, and

$$|F_N^\#| = O(n^{2d-1}) = O(N^{2(d/m)-1}) = o(N^{2\alpha}) = o(|F_N|^2),$$

verifying (1.13). \square

Proof of Theorem 1.7. All limits in the proof are as $N \rightarrow \infty$. We begin by observing that the assumption (1.12) implies

$$\sum_{j=1}^N |F_{Nj}^*|^2 \leq \max_j |F_{Nj}^*| \sum_{j=1}^N |F_{Nj}^*| \leq \max_j |F_{Nj}^*| \cdot d|F_N| = o(|F_N|^2). \quad (3.4)$$

We use the martingale central limit theorem, as stated in [11, Corollary (2.13)]. We let

$$F_{Nj} = \{(i_1, \dots, i_d) \in F_N : i_d = j\} \subseteq F_{Nj}^*,$$

and let

$$X_{Nj} = \sum_{(i_1, \dots, i_d) \in F_{Nj}} r_{i_1} \cdots r_{i_d} = r_j \sum_{(i_1, \dots, i_d) \in F_{Nj}} r_{i_1} \cdots r_{i_{d-1}}.$$

Then

$$S_N = \sum_{j=1}^N X_{Nj}$$

and, with $\tilde{X}_{Nj} = |F_N|^{-1/2} X_{Nj}$,

$$\tilde{S}_N = \sum_{j=1}^N \tilde{X}_{Nj}.$$

Evidently, $(\tilde{X}_{Nj})_{j=1}^N$ is a martingale difference sequence, and we have $\mathbb{E} \tilde{S}_N^2 = \sum \mathbb{E} \tilde{X}_{Nj}^2 = 1$.

By [11, Corollary (2.13)], to prove $\tilde{S}_N \xrightarrow{d} N(0, 1)$ it suffices to verify the Lindeberg condition

$$\sum_{j=1}^N \mathbb{E}(\tilde{X}_{Nj}^2 \mathbf{1}[|\tilde{X}_{Nj}| > \varepsilon]) \rightarrow 0 \quad \text{for every } \varepsilon > 0, \quad (3.5)$$

together with

$$\limsup_{N \rightarrow \infty} \sum_{i \neq j} \mathbb{E}(\tilde{X}_{Ni}^2 \tilde{X}_{Nj}^2) \leq 1. \quad (3.6)$$

Since every moment of \tilde{S}_N stays bounded by (1.2), moment convergence will follow as well.

To prove (3.5) it suffices to show

$$\sum_{j=1}^N \mathbb{E} \tilde{X}_{Nj}^4 \rightarrow 0. \quad (3.7)$$

In our case, by (1.2) we note $\|\tilde{X}_{Nj}\|_4 \leq 3^{d/2} \|\tilde{X}_{Nj}\|_2$ and therefore

$$\sum_{j=1}^N \mathbb{E} \tilde{X}_{Nj}^4 \leq 3^{2d} \sum_{j=1}^N \|\tilde{X}_{Nj}\|_2^4 = 3^{2d} \sum_{j=1}^N \frac{|F_{Nj}|^2}{|F_N|^2} \leq 3^{2d} |F_N|^{-2} \sum_{j=1}^N |F_{Nj}^*|^2,$$

which by (3.4) implies (3.7).

It remains to verify (3.6). For simplicity we treat first the case $d = 2$, and will describe later the modifications needed in the general case. If $d = 2$, then

$$\mathbb{E}(X_{Ni}^2 X_{Nj}^2) = \sum_{k,l,m,n} \mathbb{E} \mathbf{1}_{F_N}(k, i) \mathbf{1}_{F_N}(l, i) \mathbf{1}_{F_N}(m, j) \mathbf{1}_{F_N}(n, j) r_k r_l r_m r_n.$$

We have, $\mathbb{E} r_k r_l r_m r_n = 0$ unless the indices k, l, m, n coincide in pairs, and obtain (overcounting the case when all four indices coincide)

$$\begin{aligned} \mathbb{E}(X_{Ni}^2 X_{Nj}^2) &\leq \sum_{k,m} \mathbf{1}_{F_N}(k, i) \mathbf{1}_{F_N}(k, i) \mathbf{1}_{F_N}(m, j) \mathbf{1}_{F_N}(m, j) \\ &\quad + 2 \sum_{k,l} \mathbf{1}_{F_N}(k, i) \mathbf{1}_{F_N}(l, i) \mathbf{1}_{F_N}(k, j) \mathbf{1}_{F_N}(l, j). \end{aligned}$$

Summing the first term on the right over all i and j , we obtain $|F_N|^2$. Therefore, to show (3.6), it suffices to verify

$$\sum_{k,l} \mathbf{1}_{F_N}(k, i) \mathbf{1}_{F_N}(l, i) \mathbf{1}_{F_N}(k, j) \mathbf{1}_{F_N}(l, j) = o(|F_N|^2). \quad (3.8)$$

The sum above equals the number of pairs $((k, i), (l, j)) \in F_N \times F_N$ such that also $((l, i), (k, j)) \in F_N \times F_N$. The number of such pairs with distinct i, j, k, l is at most $|F_N^\#|$. Further, the number of pairs $((k, i), (l, j)) \in F_N \times F_N$ where two indices are equal to some r is at most $|F_{Nr}^*|^2$. Consequently, the sum in (3.8) is at most

$$|F_N^\#| + \sum_{r=1}^N |F_{Nr}^*|^2,$$

and (3.8) follows by (1.13) and (3.4).

In the case $d \geq 2$, we similarly find that $\mathbb{E} X_{Ni}^2 X_{Nj}^2$ equals the number of quadruples I_1, I_2, I_3, I_4 of d -tuples in F_N wherein the $4d$ indices coincide in pairs, and the last index is i in I_1 and I_2 and j in I_3 and I_4 . We group such quadruples according to the positions of the pairs of coinciding elements (again overcounting in the cases with less than $2d$ distinct indices, when there are several possibilities of pairing).

To do this precisely, let $\hat{I}_k = \{1, \dots, d\} \times \{k\}$, $k = 1, 2, 3, 4$; thus, $\hat{I}_1, \hat{I}_2, \hat{I}_3, \hat{I}_4$ are four disjoint copies of $\{1, \dots, d\}$. We define a *pattern* to be a complete matching in $\hat{I}_1 \cup \hat{I}_2 \cup \hat{I}_3 \cup \hat{I}_4$, i.e. a partition of the $4d$ points into $2d$ pairs, which are regarded as the edges of a graph.

For a pattern π , any assignment of indices in $\{1, \dots, N\}$ to the $2d$ edges defines 4 d -tuples I_1, I_2, I_3, I_4 in the obvious way. Let $T_N(\pi)$ be the number of quadruples $(I_1, I_2, I_3, I_4) \in F_N^4$ generated this way, i.e., the number of all assignments such that $I_1 \in F_N, I_2 \in F_N, I_3 \in F_N, I_4 \in F_N$.

Finally, let Π' denote the set of all patterns that contain the two edges $\{(d, 1), (d, 2)\}$ and $\{(d, 3), (d, 4)\}$.

In this framework, we then observe

$$\sum_{i,j} \mathbb{E} X_{Ni}^2 X_{Nj}^2 \leq \sum_{\pi \in \Pi'} T_N(\pi).$$

We classify the patterns in Π' into three types: a pattern is of type I if all its edges are inside $\hat{I}_1 \cup \hat{I}_2$ or $\hat{I}_3 \cup \hat{I}_4$; it is of type II if it is not of type I and there are no edges connecting \hat{I}_2 and \hat{I}_3 , and type III otherwise.

First, consider a pattern π of type I. Since the d -tuples in F_N are ordered, it follows that $T_N(\pi) = 0$ unless π is the pattern with edges $\{(i, 1), (i, 2)\}$ and $\{(i, 3), (i, 4)\}$, $i = 1, \dots, d$. In this case, $I_1 = I_2$ and $I_3 = I_4$, which are arbitrary elements of F_N , and thus $T_N(\pi) = |F_N|^2$.

Because the set of patterns is finite, it suffices to show that $T_N(\pi) = o(|F_N|^2)$ for every pattern π of type II or III.

If π is of type II, then I_1 and I_4 together determine I_2 and I_3 . As in the case $d = 2$, the number of allowed pairs (I_1, I_4) with distinct indices is at most $|F_N^\#|$, and the number of pairs (I_1, I_4) with at least one common index is at most $\sum_{r=1}^N |F_{Nr}^*|^2$. Therefore $T_N(\pi) = o(|F_N|^2)$ by (1.13) and (3.4).

Finally, suppose that π is of type III. Let $\hat{I}_L = \hat{I}_1 \cup \hat{I}_2$ and $\hat{I}_R = \hat{I}_3 \cup \hat{I}_4$, and call these the left and right sides of the pattern. We further say that the points $(i, k) \in \hat{I}_L$ and $(i, k + 2) \in \hat{I}_R$ are the *mirror images* of one another.

Suppose that there are r edges between \hat{I}_L and \hat{I}_R ; call these r edges crossing, and order them (in some way). Let $t_N^L(k_1, \dots, k_r)$ be the number of ways to assign indices to the edges inside \hat{I}_L such that, with k_1, \dots, k_r assigned to the crossing edges, $I_1, I_2 \in F_N$. Similarly, let $t_N^R(k_1, \dots, k_r)$ be the corresponding number of ways to assign indices in \hat{I}_R such that $I_3, I_4 \in F_N$. Then,

$$T_N(\pi) = \sum_{k_1, \dots, k_r=1}^N t_N^L(k_1, \dots, k_r) t_N^R(k_1, \dots, k_r).$$

Further, let π' be the pattern obtained by taking the edges inside \hat{I}_L in π together with their mirror images in \hat{I}_R and the edges connecting each remaining point to its mirror image. Define π'' similarly, starting with the edges inside \hat{I}_R in π . Note that both π' and π'' are patterns of type II. Then, by the Cauchy–Schwarz inequality,

$$\begin{aligned} T_N(\pi) &= \sum_{k_1, \dots, k_r=1}^N t_N^L(k_1, \dots, k_r) t_N^R(k_1, \dots, k_r) \\ &\leq \left(\sum_{k_1, \dots, k_r=1}^N t_N^L(k_1, \dots, k_r)^2 \right)^{1/2} \left(\sum_{k_1, \dots, k_r=1}^N t_N^R(k_1, \dots, k_r)^2 \right)^{1/2} \\ &= T_N(\pi')^{1/2} T_N(\pi'')^{1/2} \\ &= o(|F_N|)^2, \end{aligned}$$

where the final estimate holds because π' and π'' are of type II.

This completes the proof of (3.6) and thus of the theorem. \square

Proof of Theorem 1.8. If \tilde{S}_N converges in distribution, then (1.2) implies that all moments converge (as remarked in the proof of Theorem 1.7). In particular, $\mu = \lim \mathbb{E} \tilde{S}_N = 0$ and $\sigma^2 = \lim \mathbb{E} \tilde{S}_N^2 = 1$; further,

$$\mathbb{E} \tilde{S}_N^4 \rightarrow \mathbb{E} \xi^4 = 3. \quad (3.9)$$

Similarly, as in the proof above, ES_N^4 equals the number of quadruples (I_1, I_2, I_3, I_4) of d -tuples in F_N such that the $4d$ indices in them coincide in pairs. To estimate this number from above, we note that the number of possibilities that I_1, I_2, I_3, I_4 can coincide in two different pairs is $3|F_N|(|F_N| - 1)$, and that each element in $F_N^\#$ contributes (at least) one more to the count. Hence,

$$|F_N|^2 \mathbb{E} \tilde{S}_N^4 = \mathbb{E} S_N^4 \geq 3|F_N|^2 - 3|F_N| + |F_N^\#|. \quad (3.10)$$

Obviously, $|F_N| \rightarrow \infty$ if $\tilde{S}_N \xrightarrow{d} N(0, 1)$. Hence, (3.9) and (3.10) imply (1.13). \square

We end this section with some counterexamples where the set F_N is close to a product set and asymptotic normality does not hold.

Example 3.2. Take $F_N = \Delta_N^d$ with $d \geq 2$. It is easily seen that (1.13) does not hold, so asymptotic normality fails by Theorem 1.8. Actually, it is easy to

see that in this case, \tilde{S}_N converges to a Hermite polynomial of degree d in a standard normal variable [14], see also [7, Section XI.1] and [2, Theorem X.26].

In particular, with $d = 2$, this example shows that Theorem 1.6(i) does not extend to $\dim\{F_N\} = 2$.

Example 3.3. Fix an integer $\ell \geq 1$ and let, for $N > \ell$, F_N be the product set $\{1, \dots, \ell\} \times \{\ell + 1, \dots, N\}$.

Clearly, $S_N = \sum_1^\ell r_i \cdot \sum_{\ell+1}^N r_j$ and it follows from the central limit theorem that

$$\tilde{S}_N \xrightarrow{d} Y\xi,$$

where Y and ξ are independent, $\xi \sim N(0, 1)$ and $Y = \ell^{-1/2} \sum_1^\ell r_i$.

Hence, if $\ell = 1$, the limit is normal, but not if $\ell \geq 2$. For example, if $\ell = 2$, the limit variable is 0 with probability 1/2. (The limit can be regarded as a mixture of normal distributions with different variances.)

In particular, this example shows that Theorem 1.6(i) does not extend to $\dim\{F_N\} = 1$.

Example 3.4. Consider a *disconnected* fractional Cartesian product. For example, take $d = 6$, $m = 2$ and let S_1, \dots, S_6 be the sets $\{1, 2\}$, $\{1, 3\}$, $\{2, 3\}$, $\{4, 5\}$, $\{4, 6\}$, $\{5, 6\}$. It is easily seen that (1.13) does not hold, so asymptotic normality fails by Theorem 1.8.

This case is related to the case of disconnected G or H , respectively, in [5, Theorem 1] or [6, Theorem 1]. We expect that, as in those results, \tilde{S}_N converges to a polynomial in normal variables, but we have not checked the details.

4. FURTHER REMARKS AND OPEN PROBLEMS

Remark 4.1. It would be interesting to know more about $\|\tilde{S}_N\|_q$ and $\mathbb{P}(|\tilde{S}_N| > x)$ as functions of N . For example, how fast is the transition from the maxima in Theorem 1.5 to the limits in Theorem 1.6 as N grows?

Remark 4.2. We considered for simplicity only $a_{i_1 \dots i_d} = 1$ in Theorems 1.5 and 1.6. The upper bounds in (1.7) and (1.8) are given for arbitrary $a_{i_1 \dots i_d}$, and, in particular, $a_{i_1 \dots i_d} = \pm 1$, but the proof of the lower bounds uses the fact that all coefficients have the same sign. In general, there will be cancellations among the terms in (1.6), for any values of r_1, \dots, r_N , and it seems likely that the lower bounds in Theorem 1.5 do not extend to general $a_{i_1 \dots i_d}$. What is the correct result? Give an extension of Theorem 1.5 to arbitrary $a_{i_1 \dots i_d}$!

Certainly, the central limit theorems 1.6 and 1.7 extend to sums (1.6) with suitable conditions on $a_{i_1 \dots i_d}$, but we have not worked out the details of such extensions.

Remark 4.3. Note that if S is given by (1.6) and $d_F(\alpha) < \infty$, then

$$\|S\|_\infty \geq c \| \{a_{i_1 \dots i_d}\} \|_{\ell^{2\alpha/(\alpha+1)}},$$

where the exponent $2\alpha/(\alpha + 1)$ is the best possible; see [2, Section XIII.7]. This generalizes a result for $F = \Delta^d$ (i.e. sums (1.1), with $\alpha = d$) proved by

Littlewood [10] for $d = 2$ and for general d by [4] and [8]. It would be interesting to obtain lower bounds to the probability $\mathbb{P}(S \geq c \|\{a_{i_1 \dots i_d}\}\|_{\ell^{2\alpha/(\alpha+1)}})$.

Remark 4.4. The proofs above show that the tail estimates in Theorem 1.5 hold for the upper tails $\mathbb{P}(\tilde{S}_N > x)$ too. If d is odd, we obtain the same results for $\mathbb{P}(\tilde{S}_N < -x)$ by symmetry, but if d is even this fails. It seems likely that $\sup_N \mathbb{P}(\tilde{S}_N < -x)$ is smaller than $\exp(-cx^{2/\alpha})$ for even d , for example for $d = 2$. How small is it?

We can also replace the Rademacher system by other orthogonal systems. (See e.g. [9, Chapter 6] for a general background.)

Remark 4.5. If we replace the Rademacher variables r_i by Steinhaus functions χ_i , i.e. independent complex random variables that are uniformly distributed on the unit circle, then Theorem 1.5 still holds.

Indeed, (1.7) is still valid [2, Corollary XIII.8.29], and thus (1.8) holds by the same proof, so the upper bounds in Theorem 1.5 hold. For the lower bounds, we use the same proof as above, now taking $\mathcal{E}_N = \{\operatorname{Re} \chi_k \geq 1/2, k = 1, \dots, N\}$.

For the upper bound in (1.7), we can alternatively introduce a Rademacher system $\{r_i\}$ independent of $\{\chi_i\}$, replace χ_i by $\chi_i r_i$, which has the same distribution, and use the Rademacher version above conditioning on $\{\chi_i\}$. This standard trick works for all i.i.d. sequences of bounded symmetric random variables.

Are the central limit theorems 1.6 and 1.7 true for the Steinhaus system too, now with complex Gaussian limits? (We believe so, but we have not checked the details.)

Remark 4.6. Let us instead consider a Gaussian chaos, obtained by replacing r_i by independent Gaussian variables $\xi_i \sim N(0, 1)$.

The hypercontractive inequality (1.2) holds in this case too [12], see also [1, 7, 9], but the combinatorial dimension version (1.7) fails in the Gaussian case, as is seen by taking F to be a set with a single element.

Hence Theorem 1.5 is not true in the Gaussian case. What is true? There is no problem with the lower bounds in Theorem 1.5; the proof in Section 2 works if we take $\mathcal{E}_N = \{\xi_i > 1, i = 1, \dots, N\}$.

We believe that Theorems 1.6 and 1.7 hold for the Gaussian case too, but we have not checked the details.

Remark 4.7. Are the results true if we replace r_k by a lacunary sequence $\exp(2\pi i n_k t)$, where $\inf n_{k+1}/n_k > 1$?

5. POLARIZATION IN [2]: CORRIGENDUM

In this section we correct an argument in the proofs of Theorem VII.32 and Corollary XIII.29 in [2], results that are prominently used in the present paper. We correct also the misstatement of Theorem VII.26 ii, which led to the flawed argument.

References below are to [2]. We assume familiarity with the material in Ch. VII §8, §9, and Ch. XIII §8, and, specifically, with the notation therein.

The key error is statement (8.34) on p.165. To correct it, replace the product of the φ s on the right side of (8.34) by its symmetrization $\frac{1}{n!} \sum_{\tau \in \text{per}[n]} \varphi_{\tau 1}(j_1) \cdots \varphi_{\tau n}(j_n)$; that is, replace (8.34) by

$$\hat{\beta}_f(\varphi_1, \dots, \varphi_n) = \sum_{j_1 < \dots < j_n} \hat{f}(r_{j_1} \cdots r_{j_n}) \sum_{\tau \in \text{per}[n]} \varphi_{\tau 1}(j_1) \cdots \varphi_{\tau n}(j_n). \quad (8.34)$$

Below we list the effects of this correction.

1. In Theorem VII.26 ii replace the projective tensor algebra V_n by the algebra of the symmetric elements in V_n ; that is, replace V_n by

$$V_{n\sigma} := \{\varphi \in V_n : \varphi(j_1, \dots, j_n) = \varphi(j_{\tau 1}, \dots, j_{\tau n}) \text{ for all } \tau \in \text{per}[n]\}.$$

2. In Lemma VII.29 (p.164) replace (8.32) by

$$\hat{\mu}(r_{j_1} \cdots r_{j_n}) = \frac{1}{n!} \sum_{\tau \in \text{per}[n]} \varphi_{\tau 1}(j_1) \cdots \varphi_{\tau n}(j_n), \quad r_{j_1} \cdots r_{j_n} \in \mathbb{R}. \quad (8.32)$$

In the proof of Lemma VII.29 (p.165), on the left side of (8.36) and the left side of (8.38) replace the product of φ s by its symmetrization $\frac{1}{n!} \sum_{\tau \in \text{per}[n]} \varphi_{\tau 1}(j_1) \cdots \varphi_{\tau n}(j_n)$ (as per (8.34) above).

3. In the proof of Theorem VII.32 (p.170), replace (9.15) by

$$f_s = \sum_{j_1 < \dots < j_n} \hat{f}(r_{i_1} \cdots r_{i_n}) \left(\sum_{\tau \in \text{per}[n]} r_{i_1}(s_{\tau 1}) \cdots r_{i_n}(s_{\tau n}) \right) r_{i_1} \cdots r_{i_n}, \quad (9.15)$$

and the right side of (9.16) by $\frac{1}{n!} \sum_{\tau \in \text{per}[n]} r_{i_1}(s_{\tau 1}) \cdots r_{i_n}(s_{\tau n})$. Accordingly, replace (9.18) by

$$f = \mathbb{E}_s(F_s \star f_s) \quad (9.18)$$

and replace (9.19), (9.20), and (9.21) with the following estimations: By (9.18), Minkowski's integral inequality, and interchange of integration,

$$\begin{aligned} \|f\|_{L^p}^p &= \mathbb{E}_\omega |\mathbb{E}_s(F_s \star f_s)|^p \leq \left(\mathbb{E}_s(\mathbb{E}_\omega |F_s \star f_s|^p)^{\frac{1}{p}} \right)^p \\ &\leq \left(\mathbb{E}_s \|F_s\|_{L^1} \|f_s\|_{L^p} \right)^p \leq (n^n/n!)^p \mathbb{E}_s \mathbb{E}_\omega |f_s|^p \leq (n^n/n!)^p \mathbb{E}_\omega \mathbb{E}_s |f_s|^p. \end{aligned}$$

(We used $\|F_s\|_{L^1} \leq n^n/n!$, which follows from the corrected estimate in (8.38)). An application of the n -dimensional Khintchin inequalities (to $\mathbb{E}_s |f_s|^p$ above) implies

$$\|f\|_{L^p}^p \leq (n^n/n!)^p p^{np/2} (n!)^{p/2} \|f\|_{L^2}^p. \quad (9.21)$$

Replace (9.22) by

$$\begin{aligned} \|f\|_{L^p} &\leq \sum_{j=0}^n \|f_j\|_{L^p} \leq \sum_{j=0}^n j^j j!^{-1/2} p^{j/2} \|f_j\|_{L^2} \leq \left(\sum_{j=0}^n \frac{j^{2j} p^j}{j!} \right)^{1/2} \|f\|_{L^2} \\ &\leq e^{n/2} n^{n/2} p^{n/2} \|f\|_{L^2}. \end{aligned} \quad (9.22)$$

Following these corrections, on the right side of (9.12) replace $2e^n p^{n/2}$ by $(en)^{n/2} p^{n/2}$.

Remark 5.1. The dependence of constants on n in (9.12) is a result of the symmetrization procedure that sets up passage from the n -dimensional Khintchin inequalities to the Bonami inequalities. Bonami's direct arguments in [3] yield better constants; cf. (1.2). The same symmetrization argument is used again in Chapter XIII to affect a passage from the Khintchin inequalities in fractional dimensions to the Bonami inequalities in fractional dimensions.

4. In (12.16) on p.194, replace V_n by $V_{n\sigma}$ (algebra of symmetric elements in V_n).

5. On p.499, in (8.24) replace the right hand side by

$$\sum_{\gamma_{j_1} \cdots \gamma_{j_n} \in F} \hat{f}(\gamma_{j_1} \cdots \gamma_{j_n}) \left(\sum_{\tau \in \text{per}[n]} \gamma_{j_1}(x_{\tau 1}) \cdots \gamma_{j_n}(x_{\tau n}) \right) \gamma_{j_1} \cdots \gamma_{j_n}.$$

In (8.25) replace the right side of the first line by $\frac{1}{n!} \sum_{\tau \in \text{per}[n]} \bar{\gamma}_{j_1}(x_{\tau 1}) \cdots \bar{\gamma}_{j_n}(x_{\tau n})$, and the right side of the second line by $(2e)^n$. Continue as in 3 above, making the appropriate numerical adjustments. In particular, in (8.26) replace 64^n by $2^n e^{2n} (n!)^{1/2}$.

Acknowledgement. This research was performed while the authors visited the Mittag-Leffler Institute in Djursholm, Sweden.

REFERENCES

- [1] W. Beckner, Inequalities in Fourier analysis. *Ann. Math.* **102** (1975), 159–182.
- [2] R. Blei, *Analysis in Integer and Fractional Dimension*. Cambridge Univ. Press, Cambridge, UK, 2001.
- [3] A. Bonami, Étude des coefficients de Fourier des fonctions de $L^p(G)$. *Ann. Inst. Fourier (Grenoble)* **20** (1970), no. 2, 335–402.
- [4] A.M. Davie, Quotient algebras of uniform algebras. *J. London Math. Soc.* (2) **7** (1973), 31–40.
- [5] S. Janson, A functional limit theorem for random graphs with applications to subgraph count statistics. *Random Struct. Alg.* **1** (1990), no. 1, 15–37.
- [6] S. Janson *Orthogonal Decompositions and Functional Limit Theorems for Random Graph Statistics*. Memoirs Amer. Math. Soc. **534**, Amer. Math. Soc., Providence, R.I., 1994.
- [7] S. Janson, *Gaussian Hilbert Spaces*, Cambridge Univ. Press, Cambridge, UK, 1997.
- [8] G.W. Johnson & G.S. Woodward, On p -Sidon sets. *Indiana Univ. Math. J.* **24** (1974/75), 161–167.
- [9] S. Kwapien & W.A. Woyczyński, *Random series and stochastic integrals: single and multiple*. Birkhäuser, Boston, MA, 1992.
- [10] J.E. Littlewood, On bounded bilinear forms in an infinite number of variables. *Quart. J. Math. Oxford* **1** (1930), 164–174.
- [11] D.L. McLeish, Dependent central limit theorems and invariance principles. *Ann. Probab.* **2** (1974), 620–628.
- [12] E. Nelson, The free Markov field. *J. Funct. Anal.* **12** (1973), 211–227.
- [13] H. Rademacher, Einige Sätze über Reihen von allgemeinen Orthogonalfunktionen. *Math. Ann.* **87** (1922), 112–138.
- [14] H. Rubin & R.A. Vitale, Asymptotic distribution of symmetric statistics. *Ann. Statist.* **8** (1980), no. 1, 165–170.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CONNECTICUT, U-9 STORRS, CT
06269, USA

E-mail address: `blei@math.uconn.edu`

DEPARTMENT OF MATHEMATICS, UPPSALA UNIVERSITY, PO Box 480, SE-751 06
UPPSALA, SWEDEN

E-mail address: `svante.janson@math.uu.se`

URL: `http://www.math.uu.se/~svante`