

Some remarks on the combinatorics of \mathcal{IS}_n

Svante Janson and Volodymyr Mazorchuk

Abstract

We describe the asymptotic behavior of the cardinalities of the finite symmetric inverse semigroup \mathcal{IS}_n and its endomorphism semigroup. This is applied to show that $|\mathcal{IS}_n|/|\text{End}(\mathcal{IS}_n)|$ is asymptotically 0, solving a problem of Schein and Teclezghi. We also apply our results to compute the distributions of elements from \mathcal{IS}_n with respect to certain combinatorial properties, and to compute the generating functions for $|\mathcal{IS}_n|$ and for the number of nilpotent elements in \mathcal{IS}_n .

1 Introduction

For $n \in \mathbb{N}$ let \mathcal{IS}_n denote the symmetric inverse semigroup of all partial injections on $N_n = \{1, \dots, n\}$. We refer the reader to [GM1, GM2, Li] for the details and standard notation. For $\alpha \in \mathcal{IS}_n$ we denote by $\text{dom}(\alpha)$ the *domain* of α , by $\text{im}(\alpha)$ the *range* of α , by $\text{rank}(\alpha) = |\text{dom}(\alpha)| = |\text{im}(\alpha)|$ the *rank* of α , and by $\text{def}(\alpha) = n - \text{rank}(\alpha)$ the *defect* of α . For $k = 0, 1, \dots, n$ let $R_{n,k}$ denote the cardinality of the set $\{\alpha \in \mathcal{IS}_n : \text{rank}(\alpha) = k\}$. We immediately have

$$R_{n,k} = \binom{n}{k}^2 \cdot k!, \quad |\mathcal{IS}_n| = \sum_{i=0}^n R_{n,i} = \sum_{i=0}^n \binom{n}{i}^2 \cdot i!.$$

For elements from \mathcal{IS}_n one can use their regular tableaux presentation

$$\alpha = \begin{pmatrix} i_1 & i_2 & \dots & i_k \\ j_1 & j_2 & \dots & j_k \end{pmatrix},$$

where $\text{dom}(\alpha) = \{i_1, \dots, i_k\}$ and $\text{im}(\alpha) = \{j_1, \dots, j_k\}$. However, sometimes it is more convenient to use the so-called *chain* (or *chart*) decomposition of α , which is analogous to the cyclic decomposition for usual permutations. We refer the reader to [Li] for rigorous definitions, however, this decomposition is very easy to explain on the following example. The element

$$\alpha = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 7 & 9 \\ 7 & 4 & 5 & 1 & 10 & 2 & 6 \end{pmatrix} \in \mathcal{IS}_{10}$$

has the following graph of the action on $\{1, 2, \dots, 10\}$:

$$\begin{array}{ccccccc} 1 & \rightarrow & 7 & & & & \\ \uparrow & & \downarrow & & 3 \rightarrow 5 \rightarrow 10 & & 9 \rightarrow 6 & & 8, \\ 4 & \leftarrow & 2 & & & & & & \end{array}$$

and hence it is convenient to write it as $\alpha = (1, 7, 2, 4)[3, 5, 10][9, 6][8]$. We call $(1, 7, 2, 4)$ a *cycle* and $[3, 5, 10]$ (as well as $[9, 6]$ and $[8]$) a *chain* of the element α .

We denote by L_n the total number of chains in the chain decompositions of all elements in $|\mathcal{IS}_n|$. Each element of rank k has defect $n - k$ and thus contains $n - k$ chains implying $L_n = \sum_{k=0}^n (n - k)R_{n,k}$. The semigroup \mathcal{IS}_n contains the zero element 0 , uniquely characterized by the property $\text{dom}(0) = \emptyset$. We denote by T_n the set of all nilpotent elements in \mathcal{IS}_n , that is the set of all $\alpha \in \mathcal{IS}_n$ satisfying $\alpha^n = 0$. We also denote by $L^{(n)}$ the total number of chains in the chain decompositions of all elements in T_n .

In [GM2] various combinatorial relations between $|\mathcal{IS}_n|$, $|T_n|$, L_n and $L^{(n)}$ were obtained in a purely combinatorial way. The paper [GM2] contains also various estimates of distributions of elements from \mathcal{IS}_n with respect to certain algebraic properties. These distributions are obtained using several technical lemmas. The most of the technical difficulties in [GM2] arise from the fact that the authors did not have any reasonable asymptotic formula for $|\mathcal{IS}_n|$ available. The aim of the present paper is to fill this gap. In Section 2 we derive an asymptotic formula for $|\mathcal{IS}_n|$. In Section 4 we even show that analogous methods can be applied to derive an asymptotic formula for $|\text{End}(\mathcal{IS}_n)|$. These formulae happen to be enough to show that $|\mathcal{IS}_n|/|\text{End}(\mathcal{IS}_n)| \rightarrow 0$, $n \rightarrow \infty$, which solves a problem from [ST]. Our results can be used to recover (in hopefully an easier way) several asymptotic statements from [GM2]. This is done in Section 3. Our results can be also used to obtain several new statements about the distributions of elements of \mathcal{IS}_n with respect to such combinatorial properties as the defect, the stable rank, the order etc. This is done in Section 5. Finally, in Section 6 we compute exponential generating functions for $|\mathcal{IS}_n|$, $|T_n|$, L_n and $L^{(n)}$ and use them to recover various combinatorial results from [GM2].

2 An asymptotic for $|\mathcal{IS}_n|$

This section is devoted to the proof of the following

Theorem 1.

$$|\mathcal{IS}_n| \sim \frac{1}{2\sqrt{\pi e}} n^{-1/4} e^{2\sqrt{n}} n! \sim \frac{1}{\sqrt{2e}} \cdot e^{2\sqrt{n}-n} n^{n+1/4}.$$

Proof. For $R_{n,k} = \binom{n}{k}^2 \cdot k! = \frac{n!^2}{(n-k)!^2 k!}$ we have the ratio $\frac{R_{n,k+1}}{R_{n,k}} = \frac{(n-k)^2}{k+1}$. Moreover, for large n we obtain that $\frac{R_{n,k+1}}{R_{n,k}} \approx 1$ when $k \approx n - \sqrt{n}$, hence $\max_k R_{n,k}$ is achieved for such a k . Note that $\frac{R_{n,k+1}}{R_{n,k}}$ is decreasing with respect to k . Write

$$k = n - x\sqrt{n}, \quad 0 \leq x \leq \sqrt{n}. \quad (1)$$

Using the Stirling formula we have

$$\begin{aligned} \ln \left(\frac{R_{n,k}}{n!} \right) &= n \ln n - n + \frac{1}{2} \ln(2\pi n) - k \ln k + k - \frac{1}{2} \ln(2\pi k) - \\ &\quad - 2(n-k) \ln(n-k) + 2(n-k) - \ln(2\pi(n-k)) + O \left(\frac{1}{n} + \frac{1}{k} + \frac{1}{n-k} \right). \end{aligned} \quad (2)$$

Using the arguments above we have $\frac{R_{n,k+1}}{R_{n,k}} < \frac{(\frac{1}{2}\sqrt{n})^2}{n - \frac{1}{2}\sqrt{n}} < \frac{1}{2}$ for $k > n - \frac{1}{2}\sqrt{n}$ and large n . Thus, for $k \geq k_1 = \lceil n - \frac{1}{2}\sqrt{n} \rceil$ we have $R_{n,k} \leq 2^{-(k-k_1)} R_{n,k_1}$. In particular,

$$\sum_{k \geq n - \frac{1}{4}\sqrt{n}} R_{n,k} \leq 2^{2 - \frac{1}{4}\sqrt{n}} R_{n,k_1} = O \left(2^{-\sqrt{n}/4} R_{n,k_1} \right).$$

Similarly, for $k \leq n - 2\sqrt{n}$ we have $\frac{R_{n,k}}{R_{n,k+1}} < \frac{n}{(2\sqrt{n})^2} = \frac{1}{4}$, and

$$\sum_{k \leq n - 3\sqrt{n}} R_{n,k} = O \left(4^{-\sqrt{n}} R_{n,k_2} \right),$$

where $k_2 = \lceil n - 2\sqrt{n} \rceil$.

Hence, to estimate $|\mathcal{IS}_n| = \sum_{k=0}^n R_{n,k}$ we can ignore $k \geq n - \frac{1}{4}\sqrt{n}$ and $k \leq n - 3\sqrt{n}$. We may thus assume that $\frac{1}{4} \leq x \leq 3$. For such x we have:

$$\begin{aligned} \ln \left(\frac{R_{n,k}}{n!} \right) &= n \ln n - n - (n - x\sqrt{n}) \ln(n - x\sqrt{n}) + n - x\sqrt{n} - \frac{1}{2} \ln \frac{n - x\sqrt{n}}{n} - \\ &\quad - 2x\sqrt{n} \ln x - 2x\sqrt{n} \ln(\sqrt{n}) + 2x\sqrt{n} - \ln(2\pi x\sqrt{n}) + O(n^{-1/2}) = \\ &= - (n - x\sqrt{n}) \ln \left(1 - \frac{x}{\sqrt{n}} \right) - 2\sqrt{n}x \ln x + x\sqrt{n} - \ln(2\pi x\sqrt{n}) + O(n^{-1/2}) = \\ &= x\sqrt{n} - x^2 + \frac{x^2}{2} + x\sqrt{n} - 2\sqrt{n}x \ln x - \ln(2\pi x\sqrt{n}) + O(n^{-1/2}) = \\ &= 2\sqrt{n}(x - x \ln x) - \frac{x^2}{2} - \ln x - \ln(2\pi\sqrt{n}) + O(n^{-1/2}), \end{aligned}$$

where all O are uniform in x and n .

Denote $f(x) = x - x \ln x$ and we have $f'(x) = -\ln x$, $f''(x) = -\frac{1}{x}$. Thus $f(x)$ is concave on $[0, +\infty)$ with a maximum at $x_0 = 1$. As $f(x_0) = 1$, we have the following Taylor expansion:

$$f(x) = 1 - \frac{1}{2}(x - x_0)^2 + O(|x - x_0|^3), \quad 0 \leq x < \infty. \quad (3)$$

For $\frac{1}{4} \leq x \leq 3$ we have $f''(x) < -\frac{1}{3}$ and thus $f(x) \leq 1 - \frac{1}{6}(x - x_0)^2$.

Further, let $g(x) = -\frac{x^2}{2} - \ln x$. Then for all $\frac{1}{4} \leq x \leq 3$ such that $x\sqrt{n} \in \mathbb{Z}$ we have

$$\frac{1}{n!} R_{n, n-x\sqrt{n}} = e^{2\sqrt{n}f(x)+g(x)} \cdot \frac{1 + O(n^{-1/2})}{2\pi\sqrt{n}}. \quad (4)$$

Now we have:

$$\begin{aligned} \frac{1}{n!} \sum_{k=0}^n R_{n,k} &= \int_0^{n+1} \frac{1}{n!} R_{n, n-[t]} dt \sim \int_{\sqrt{n}/4}^{3\sqrt{n}} \frac{1}{n!} R_{n, n-[t]} dt = [t = \sqrt{n}y] = \\ &= \sqrt{n} \int_{1/4}^3 \frac{1}{n!} R_{n, n-[y\sqrt{n}]} dy = \left[\tilde{y} = \frac{[y\sqrt{n}]}{\sqrt{n}} \right] = \sqrt{n} \int_{1/4}^3 \frac{1 + O(n^{-1/2})}{2\pi\sqrt{n}} e^{2\sqrt{n}f(\tilde{y})+g(\tilde{y})} dy \sim \\ &\sim \frac{e^{2\sqrt{n}}}{2\pi} \int_{1/4}^3 e^{2\sqrt{n}(f(\tilde{y})-1)+g(\tilde{y})} dy. \end{aligned}$$

Write

$$\int_{1/4}^3 e^{2\sqrt{n}(f(\tilde{y})-1)+g(\tilde{y})} dy = \int_{I_1} e^{2\sqrt{n}(f(\tilde{y})-1)+g(\tilde{y})} dy + \int_{I_2} e^{2\sqrt{n}(f(\tilde{y})-1)+g(\tilde{y})} dy,$$

where $I_1 = \{y \in [1/4, 3] : |y-1| \geq n^{-1/5}\}$ and $I_2 = \{y \in [1/4, 3] : |y-1| \leq n^{-1/5}\}$ and denote these integrals by X_1 and X_2 respectively.

Since $|\tilde{y} - y| < n^{-1/2}$, for $1/4 \leq y \leq 3$ we have

$$2\sqrt{n}(f(\tilde{y}) - 1) + g(\tilde{y}) \leq -2\sqrt{n} \frac{(\tilde{y} - 1)^2}{6} + O(1) = -\frac{\sqrt{n}}{3}(y - 1)^2 + O(1).$$

Hence $X_1 = O(e^{-n^{1/10}/3})$.

From (3) we also have, uniformly for $y \in I_2$, that

$$\begin{aligned} 2\sqrt{n}(f(\tilde{y}) - 1) &= 2\sqrt{n} \left(-\frac{1}{2}(\tilde{y} - 1)^2 + O(n^{-3/5}) \right) = -\sqrt{n}(\tilde{y} - 1)^2 + O(n^{-1/10}) = \\ &= -\sqrt{n}(y - 1)^2 + O(n^{1/2-1/5}|\tilde{y} - y| + n^{-1/10}) = -\sqrt{n}(y - 1)^2 + o(1), \end{aligned}$$

and, similarly, $g(\tilde{y}) = g(1) + O(n^{-1/5}) = -1/2 + o(1)$.

Now we calculate again:

$$\begin{aligned} \frac{1}{n!} \sum_{k=0}^n R_{n,k} &\sim \frac{e^{2\sqrt{n}}}{2\pi} (X_1 + X_2) \sim \frac{e^{2\sqrt{n}}}{2\pi} X_2 \sim \frac{e^{2\sqrt{n}}}{2\pi} \int_{1-n^{-1/5}}^{1+n^{-1/5}} e^{-\sqrt{n}(y-1)^2-1/2} dy \sim \\ &\sim \frac{e^{2\sqrt{n}-1/2}}{2\pi} \int_{-\infty}^{+\infty} e^{-\sqrt{n}(y-1)^2} dy = \frac{e^{2\sqrt{n}-1/2}}{2\pi} \sqrt{\frac{\pi}{\sqrt{n}}} = \frac{1}{2} \pi^{-1/2} e^{-1/2} n^{-1/4} e^{2\sqrt{n}}. \end{aligned}$$

Finally, using the Stirling formula again, we obtain

$$|\mathcal{IS}_n| = \sum_{k=0}^n R_{n,k} \sim \frac{1}{2\sqrt{\pi}e} n^{-1/4} e^{2\sqrt{n}} n! \sim \frac{1}{\sqrt{2}e} n^{n+1/4} e^{2\sqrt{n}-n},$$

completing the proof. \square

3 Some applications of Theorem 1

An immediate corollary of Theorem 1 is the following statement, proved in [GM2, Theorem 8]:

Corollary 1.

$$\frac{|\mathcal{IS}_{n+1}|}{|\mathcal{IS}_n|} \sim n, \quad n \rightarrow \infty.$$

Another corollary is the following reinforcement of [GM2, Theorem 9]:

Corollary 2. $|T_n| \sim \frac{1}{\sqrt{n}}|\mathcal{IS}_n|$, in particular,

$$\frac{|T_n|}{|\mathcal{IS}_n|} \rightarrow 0, \quad n \rightarrow \infty.$$

Proof. From [GM2, Theorem 6] we know that $|T_n| = \frac{1}{n}L_n$ (see a different proof in Section 6). By the definition, $L_n = \sum_{k=0}^n (n-k)R_{n,k}$. An argument, analogous to that of Theorem 1, yields

$$\frac{1}{n!}L_n \sim \int_{\sqrt{n}/4}^{3\sqrt{n}} [t] \frac{1}{n!} R_{n,n-[t]} dt.$$

The same estimates as in Theorem 1 show that most of the integral comes from $y = t/\sqrt{n} = 1 + O(n^{-1/5})$. Hence

$$\frac{1}{n!}L_n \sim \sqrt{n} \int_{\sqrt{n}/4}^{3\sqrt{n}} \frac{1}{n!} R_{n,n-[t]} dt \sim \sqrt{n} \frac{1}{n!} |\mathcal{IS}_n|.$$

This implies that $L_n \sim \sqrt{n}|\mathcal{IS}_n|$ and completes the proof. \square

4 An asymptotic for $|\text{End}(\mathcal{IS}_n)|$

In [ST] it is shown that for $n > 6$ the cardinality of the semigroup $\text{End}(\mathcal{IS}_n)$ of all endomorphisms of the semigroup \mathcal{IS}_n equals

$$|\text{End}(\mathcal{IS}_n)| = 3^n + 3 \cdot n! + n! \sum_{m=0}^n \sum_{k=1}^{\lfloor m/2 \rfloor} \frac{2^{m-3k}}{(n-m)! \cdot (m-2k)! \cdot k!}.$$

On [ST, Page 303] the following problem is formulated:

Find an asymptotic estimate for $|\text{End}(\mathcal{IS}_n)|$ when $n \rightarrow \infty$. Is $|\text{End}(\mathcal{IS}_n)|/|\mathcal{IS}_n|$ approaching 0?

In this section we answer both parts of this problem.

Theorem 2. $|\text{End}(\mathcal{IS}_n)| \sim 3n!$.

Proof. Set

$$X_n = n! \sum_{m=0}^n \sum_{k=1}^{\lfloor m/2 \rfloor} \frac{2^{m-3k}}{(n-m)! \cdot (m-2k)! \cdot k!}.$$

It would be enough to show that $X_n/n! \rightarrow 0$, $n \rightarrow \infty$. To do this we remark that X_n equals the number of ways to perform the following procedure:

- (i) choose $X \subset N_n$;
- (ii) choose $Y \subset X$ such that $|Y| = 2k > 0$;
- (iii) decompose $Y = \cup Y_i$, $|Y_i| = 2$, $Y_i \cap Y_j = \emptyset$ for $i \neq j$, the order of Y_i is not important;
- (iv) Choose $Z \subset X \setminus Y$.

Now let $|X| = m$, $0 \leq m \leq n$, and note that (i) can be done in $\binom{n}{m}$ different ways, each of (ii) and (iv) can be done in at most 2^m different ways, and, finally, (iii) can be done in at most $m!!$ different ways. Hence

$$X_n \leq \sum_{m=0}^n \binom{n}{m} \cdot 2^m \cdot 2^m \cdot m!! \leq \left(\sum_{m=0}^n \binom{n}{m} \cdot 4^m \right) (2^{\lceil n/2 \rceil})!! = 5^n 2^{\lceil n/2 \rceil} \lceil n/2 \rceil!.$$

To complete the proof it is enough to show that $5^n 2^{\lceil n/2 \rceil} \lceil n/2 \rceil! / n! \rightarrow 0$, $n \rightarrow \infty$. Using the Stirling formula we have

$$5^n 2^{\lceil n/2 \rceil} \lceil n/2 \rceil! \leq 5^n 2^{(n+1)/2} \lceil n/2 \rceil! \sim \frac{1}{\sqrt{\pi}} e^{n \ln 5\sqrt{2} - \frac{1}{2} \ln \lceil n/2 \rceil + \lceil n/2 \rceil \ln \lceil n/2 \rceil - \lceil n/2 \rceil},$$

and thus

$$\frac{5^n 2^{\lceil n/2 \rceil} \lceil n/2 \rceil!}{n!} \sim \frac{1}{\sqrt{2}} e^{n \ln 5\sqrt{2} - \frac{1}{2} \ln \lceil n/2 \rceil + \lceil n/2 \rceil \ln \lceil n/2 \rceil - \lceil n/2 \rceil - \frac{1}{2} \ln n - n \ln n}.$$

Since the exponent is $-\frac{1}{2}n \ln n + O(n)$, we obtain that the expression approaches 0 for large n . This completes the proof. \square

Corollary 3.

$$\frac{|\text{End}(\mathcal{IS}_n)|}{|\mathcal{IS}_n|} \rightarrow 0, \quad n \rightarrow \infty.$$

Proof. Follows immediately from the formulae of Theorem 1 and Theorem 2. \square

Using the methods, analogous to those of Theorem 1, one can even estimate the asymptotic for the ‘‘problematic’’ term X_n above.

Theorem 3.

$$X_n \sim \frac{1}{\sqrt{2}} \cdot e^{\frac{1}{2}n \ln n - \frac{1}{2}n + 3\sqrt{n} - \frac{9}{4}}.$$

Proof. We can write

$$n! \sum_{m=0}^n \sum_{k=1}^{\lfloor m/2 \rfloor} \frac{2^{m-3k}}{(n-m)! \cdot (m-2k)! \cdot k!} = n! \sum_{m=0}^n \frac{2^m}{(n-m)!} \sum_{k=1}^{\lfloor m/2 \rfloor} \frac{2^{-3k}}{(m-2k)! \cdot k!},$$

and we denote $a_k = \frac{2^{-3k}}{(m-2k)! \cdot k!}$, $b_m = \sum_{k=1}^{\lfloor m/2 \rfloor} a_k$, $c_m = \frac{2^m}{(n-m)!} b_m$. Remark that $\frac{a_{k+1}}{a_k} = \frac{2^{-3}}{k+1} (m-2k)(m-2k-1)$ decreases on $[0, m/2)$ and a_k has on $[0, m/2)$ a unique maximum at $\approx \frac{m}{2} - \sqrt{m}$. Let $k = \frac{m}{2} - x\sqrt{m}$, that is $m-2k = 2x\sqrt{m}$, where $\varepsilon \leq x \leq C$. Then we have

$$\frac{a_{k+1}}{a_k} = \frac{1}{8} \cdot \frac{1}{m/2} \cdot 4x^2 m (1 + o(1)) = x^2 + o(1).$$

This implies that $\sum_{x < 1/2} a_k$ and $\sum_{x > 2} a_k$ belong to $O(e^{-c\sqrt{m}} a_{m/2 - \sqrt{m}})$, that is relatively very small and can be neglected. Assume now that $1/2 \leq x \leq 2$. Taking into account that

$$\ln k = \ln \frac{m}{2} + \ln \left(1 - \frac{2x}{\sqrt{m}} \right) = \ln \frac{m}{2} - \frac{2x}{\sqrt{m}} - \frac{2x^2}{m} + O(m^{-3/2})$$

and using the Stirling formula we obtain the following:

$$\begin{aligned} \ln a_k &= -3k \ln 2 - \ln(2x\sqrt{m})! - \ln k! = -\frac{3 \ln 2}{2} m + 3 \ln 2\sqrt{m}x - 2x\sqrt{m} \ln 2 - \\ &- 2x\sqrt{m} \ln x - x\sqrt{m} \ln m + 2x\sqrt{m} - \ln(2\pi) - \frac{1}{2} \ln(2x\sqrt{m}) - k \ln k + k - \frac{1}{2} \ln k + o(1) = \\ &= -\frac{3 \ln 2}{2} m + (\ln 2 + 2)x\sqrt{m} - 2x\sqrt{m} \ln x - x\sqrt{m} \ln m - \ln(2\pi) - \frac{1}{2} \ln(2x\sqrt{m}) - \\ &\quad - k \ln \frac{m}{2} + \frac{2kx}{\sqrt{m}} + \frac{mx^2}{m} + \frac{m}{2} - x\sqrt{m} - \frac{1}{2} \ln \frac{m}{2} + o(1) = \\ &= -m \ln 2 + 2x\sqrt{m} - 2x\sqrt{m} \ln x - \frac{1}{2} \ln(4\pi^2 x m^{3/2}) - \frac{1}{2} m \ln m - x^2 + \frac{m}{2} + o(1). \end{aligned}$$

Further, assuming $x = 1 + m^{-1/4}y$ yields $x \ln x - x = -1 + \frac{1}{2}(x-1)^2 + O((x-1)^3) = -1 + \frac{y^2}{2\sqrt{m}} + O\left(\frac{y^3}{m^{3/4}}\right)$ and thus

$$\ln a_k = -\frac{1}{2} m \ln m + m \left(\frac{1}{2} - \ln 2 \right) - 1 + O(y m^{-1/4}) + 2\sqrt{m} - y^2 + O(y^3 m^{-1/4}) - \frac{3}{4} \ln m - \ln(2\pi).$$

Therefore $k = \frac{m}{2} - \sqrt{m} - m^{1/4}y$ yields

$$a_k = \frac{1}{2\pi} e^{(\frac{1}{2} - \ln 2)m - \frac{1}{2} m \ln m - \frac{3}{4} \ln m + 2\sqrt{m} - 1} e^{-y^2} \left(1 + O\left(\frac{y + y^3}{m^{1/4}}\right) \right).$$

We can assume that, say, $y = O(m^{1/12})$ and ignore larger y . In this way we obtain

$$\begin{aligned} b_m &= \sum_{k=1}^{\lfloor m/2 \rfloor} a_k = \frac{1}{2\pi} e^{(\frac{1}{2}-\ln 2)m - \frac{1}{2}m \ln m - \frac{3}{4} \ln m + 2\sqrt{m}-1} m^{1/4} \int_{-\infty}^{\infty} e^{-y^2} dy (1 + o(1)) \sim \\ &\sim \frac{1}{2\sqrt{\pi}} e^{(\frac{1}{2}-\ln 2)m - \frac{1}{2}m \ln m - \frac{1}{2} \ln m + 2\sqrt{m}-1}. \end{aligned}$$

The latter implies

$$\ln b_m = \left(\frac{1}{2} - \ln 2 \right) m - \frac{1}{2} m \ln m - \frac{1}{2} \ln m + 2\sqrt{m} - 1 - \ln(2\sqrt{\pi}) + o(1) \quad (5)$$

and also

$$\ln c_m = \ln b_m + m \ln 2 - \ln((n-m)!). \quad (6)$$

Further, for $m \rightarrow \infty$ we compute $\ln \frac{b_{m+1}}{b_m} = \frac{1}{2} - \ln 2 - \frac{1}{2} \ln m - \frac{1}{2} + o(1) = -\ln 2 - \frac{1}{2} \ln m + o(1)$ and also $\ln \frac{c_{m+1}}{c_m} = -\frac{1}{2} \ln m + \ln(n-m) + o(1)$. This gives us that for large n the value of c_m is largest when $\frac{1}{2} \ln m \approx \ln(n-m)$ that is $m \approx n - \sqrt{n}$. In particular, it follows easily that $m \leq n/2$ can be ignored and thus we obtain that $o(1)$, $m \rightarrow \infty$, is small even if $n \rightarrow \infty$.

Let us now show that even $m < n - 3\sqrt{n}$ can be ignored. If $m < n - 2\sqrt{n}$ then we have $-\frac{1}{2} \ln m + \ln(n-m) > -\frac{1}{2} \ln n + \ln(2\sqrt{n}) = \ln 2$ and thus for large n we derive $\ln \frac{c_{m+1}}{c_m} > 1/2$ and hence $\frac{c_{m+1}}{c_m} > e^{1/2}$. Set $M = \lceil n - 2\sqrt{n} \rceil$. Then $\frac{c_m}{c_M} < e^{-(M-m)/2}$ and thus

$$\sum_{m < n - 3\sqrt{n}} c_m < e^{-\sqrt{n}/2} \frac{1}{1 - e^{-1/2}} c_M.$$

The latter implies that all terms with $m < n - 3\sqrt{n}$ can be ignored. Similarly, all terms with $m > n - \sqrt{n}/2$ can be ignored.

Thus we can assume $m = n - x\sqrt{n}$, where $1/2 \leq x \leq 3$. Under such assumption we have $\ln \frac{c_{m+1}}{c_m} = -\frac{1}{2} \ln n + \ln(x\sqrt{n}) + o(1) = \ln x + o(1)$.

For $1/2 \leq x \leq 3$ we have, using the Stirling formula, that

$$\begin{aligned} \ln m &= \ln n + \ln \left(1 - \frac{x}{\sqrt{n}} \right) = \ln n - \frac{x}{\sqrt{n}} - \frac{x^2}{2n} + O(n^{-3/2}), \\ m \ln m &= n \ln n - x\sqrt{n} \ln n - x\sqrt{n} + x^2 - \frac{x^2}{2} + O(n^{-1/2}), \\ \ln(n-m)! &= \ln(x\sqrt{n})! = x\sqrt{n} \ln x + \frac{1}{2} x\sqrt{n} \ln n - x\sqrt{n} + \frac{1}{2} \ln x + \frac{1}{4} \ln n + \ln \sqrt{2\pi} + o(1), \\ \sqrt{m} &= \sqrt{n} (1 - x/\sqrt{n})^{1/2} = \sqrt{n} - x/2 + o(1). \end{aligned}$$

Hence, using (5) and (6), we obtain

$$\begin{aligned}\ln c_m &= \frac{1}{2}n - \frac{1}{2}x\sqrt{n} - \frac{1}{2}n \ln n + \frac{1}{2}x\sqrt{n} \ln n + \frac{1}{2}x\sqrt{n} - \frac{x^2}{4} - \frac{1}{2} \ln n + 2\sqrt{n} - x - 1 - \ln(2\sqrt{\pi}) - \\ &\quad - x\sqrt{n} \ln x - \frac{1}{2}x\sqrt{n} \ln n + x\sqrt{n} - \frac{1}{2} \ln x - \frac{1}{4} \ln n - \ln \sqrt{2\pi} + o(1) = \\ &= \frac{1}{2}n - \frac{1}{2}n \ln n - \frac{3}{4} \ln n + 2\sqrt{n} - 1 - \ln(2^{3/2}\pi) + \sqrt{n}(x - x \ln x) - \frac{x^2}{4} - x - \frac{1}{2} \ln x + o(1).\end{aligned}$$

Setting $x = 1 + yn^{-1/4}$ yields

$$\ln c_m = \frac{1}{2}n - \frac{1}{2}n \ln n - \frac{3}{4} \ln n + 3\sqrt{n} - 1 - \ln(2^{3/2}\pi) - \frac{5}{4} - \frac{y^2}{2} + O\left(\frac{y^3}{n^{1/4}}\right) + o(1)$$

and thus

$$\begin{aligned}\sum_{m=0}^n c_m &= \exp\left(\frac{1}{2}n - \frac{1}{2}n \ln n - \frac{3}{4} \ln n + 3\sqrt{n} - \frac{9}{4} - \ln(2^{3/2}\pi)\right) n^{1/4} \times \\ &\quad \times \int_{-\infty}^{+\infty} e^{-y^2/2} dy (1 + o(1)) = \frac{1}{2\sqrt{\pi}} e^{\frac{1}{2}n - \frac{1}{2}n \ln n - \frac{1}{2} \ln n + 3\sqrt{n} - \frac{9}{4} + o(1)}.\end{aligned}$$

This implies that

$$X_n \sim \frac{n!}{2\sqrt{\pi n}} e^{\frac{1}{2}n - \frac{1}{2}n \ln n + 3\sqrt{n} - \frac{9}{4}} \sim \frac{1}{\sqrt{2}} e^{\frac{1}{2}n \ln n - \frac{1}{2}n + 3\sqrt{n} - \frac{9}{4}},$$

and completes the proof. \square

5 Some distributions

Denote by D_n the defect of a random element of \mathcal{IS}_n , by X_n the stable rank of a random element of \mathcal{IS}_n , by C_n the number of cycles of a random element of \mathcal{IS}_n , and by $K_n = C_n + D_n$ the total number of components (i.e. cycles and chains) of a random element of \mathcal{IS}_n .

Proposition 1. *If $n \rightarrow \infty$ and $\frac{k-\sqrt{n}}{n^{1/4}} \rightarrow z$ with $-\infty < z < \infty$, then*

$$P(D_n = k) \sim \frac{1}{\sqrt{\pi n^{1/4}}} e^{-z^2}.$$

In particular,

$$\frac{D_n - \sqrt{n}}{n^{1/4}} \xrightarrow{d} N(0, 1/2).$$

Proof. We have $P(D_n = k) = \frac{R_{n,n-k}}{|\mathcal{IS}_n|}$ by definition and $\frac{R_{n,n-k}}{|\mathcal{IS}_n|} \sim \frac{1}{\sqrt{\pi n^{1/4}}} e^{-z^2}$, follows from (3) and (4). \square

Proposition 2.

$$P(X_n = k) \sim \frac{1}{\sqrt{n}} e^{-k/\sqrt{n}} \text{ if } k = o(n^{3/4}),$$

in particular,

$$\frac{X_n}{\sqrt{n}} \xrightarrow{d} \exp(1).$$

Proof. We have

$$P(X_n = k) = \binom{n}{k} \cdot k! \cdot \frac{|T_{n-k}|}{|\mathcal{IS}_n|} = \frac{|T_{n-k}|/(n-k)!}{|\mathcal{IS}_n|/n!}.$$

Hence, if $k = o(n)$ we have, using Section 2 and Section 3, that

$$\begin{aligned} P(X_n = k) &\sim \frac{(n-k)^{-3/4} e^{2\sqrt{n-k}}}{n^{-1/4} e^{2\sqrt{n}}} \sim \frac{1}{\sqrt{n}} e^{2(\sqrt{n-k}-\sqrt{n})} = \\ &= \frac{1}{\sqrt{n}} e^{2\sqrt{n}((1-k/n)^{1/2}-1)} = \frac{1}{\sqrt{n}} e^{-2\sqrt{n} \cdot \frac{k}{2n} + O(k^2/n^{3/2})} \end{aligned}$$

and the statement follows. \square

Proposition 3.

$$\frac{C_n - \frac{1}{2} \ln n}{\sqrt{\frac{1}{2} \ln n}} \xrightarrow{d} N(0, 1).$$

Proof. Given X_n , the number of cycles for the permutational part of size X_n is approximately $\ln X_n$. More precisely, by [Go], we have

$$\frac{C_n - \ln X_n}{\sqrt{\ln X_n}} \xrightarrow{d} N(0, 1).$$

Further, we have $\ln X_n = \frac{1}{2} \ln n + \ln \frac{X_n}{\sqrt{n}}$ and $\ln \frac{X_n}{\sqrt{n}} \xrightarrow{d} \ln \exp(1)$ by Proposition 2. Hence, in

$$\frac{C_n - \frac{1}{2} \ln n}{\sqrt{\frac{1}{2} \ln n}} = \frac{\sqrt{\ln X_n}}{\sqrt{\frac{1}{2} \ln n}} \cdot \frac{C_n - \ln X_n}{\sqrt{\ln X_n}} + \frac{\ln X_n - \frac{1}{2} \ln n}{\sqrt{\frac{1}{2} \ln n}}$$

we have $\frac{\sqrt{\ln X_n}}{\sqrt{\frac{1}{2} \ln n}} \xrightarrow{p} 1$ and $\frac{\ln X_n - \frac{1}{2} \ln n}{\sqrt{\frac{1}{2} \ln n}} \xrightarrow{p} 0$, completing the proof. \square

More precisely, we can show that C_n is almost Poisson distributed. Let d_{TV} denote the *total variation distance* between two distributions, see e.g. [BHJ].

Proposition 4.

$$d_{TV} \left(C_n, \text{Po} \left(\frac{1}{2} \ln n \right) \right) \rightarrow 0, \quad n \rightarrow \infty.$$

Proof. Let $h_k = \sum_{i=1}^k 1/i = \ln k + O(1)$. Given $X_n = k$, the number of cycles is distributed as the number of cycles in a random permutation of length k . Using [BH], we obtain

$$d_{TV}(\mathcal{L}(C_n|X_n = k), \text{Po}(h_k)) \leq \frac{c}{h_k} \leq \frac{c}{\ln k}$$

for some constant $c \leq \pi^2/6$. Further, by [BHJ, Remark 1.1.4], we have

$$d_{TV}(\text{Po}(h_k), \text{Po}(\ln \sqrt{n})) \leq \frac{|h_k - \ln \sqrt{n}|}{\sqrt{\ln \sqrt{n}}} \leq \frac{|\ln k - \ln \sqrt{n}| + 1}{\sqrt{\ln \sqrt{n}}}.$$

Consequently,

$$d_{TV}(\mathcal{L}(C_n|X_n = k), \text{Po}(\ln \sqrt{n})) \leq f(k) := \frac{\pi^2}{6 \ln k} + \frac{|\ln k - \ln \sqrt{n}| + 1}{\sqrt{\ln \sqrt{n}}}.$$

Since also $d_{TV} \leq 1$, we obtain $d_{TV}(C_n, \text{Po}(\ln \sqrt{n})) \leq E(f(X_n) \wedge 1)$. From the proof of Proposition 3 it follows that $f(X_n) \xrightarrow{p} 0$ and thus $E(f(X_n) \wedge 1) \xrightarrow{p} 0$, completing the proof. \square

Corollary 4.

$$\frac{K_n - \sqrt{n}}{n^{1/4}} \xrightarrow{d} N(0, 1/2).$$

Proof. Follows from Propositions 1 and 3. \square

Recall that for $\sigma \in \mathcal{IS}_n$ the *order* $O(\sigma)$ of σ is defined as the cardinality of the monoid, generated by σ , and the *inverse order* $\text{IO}(\sigma)$ of σ is defined as the cardinality of the inverse monoid, generated by σ , that is

$$O(\sigma) = |\{\sigma^l : l \in \{0, 1, 2, \dots\}\}|, \quad \text{IO}(\sigma) = |\{\sigma^l : l \in \mathbb{Z}\}|.$$

Let O_n and I_n denote the order and the inverse order of a random element of \mathcal{IS}_n respectively.

Proposition 5.

$$\frac{\ln O_n - \frac{1}{8} \ln^2 n}{\sqrt{\frac{1}{24} \ln^3 n}} \xrightarrow{d} N(0, 1), \quad \frac{\ln I_n - \frac{1}{8} \ln^2 n}{\sqrt{\frac{1}{24} \ln^3 n}} \xrightarrow{d} N(0, 1).$$

Proof. For $\sigma \in \mathcal{IS}_n$ denote $X(\sigma) = \{i \in \{1, \dots, n\} : \sigma^l(i) = i \text{ for some } l > 0\}$. Then $|X(\sigma)|$ is the stable rank of σ . Moreover, any $\sigma \in \mathcal{IS}_n$ can be written as a product $\sigma = \sigma_1 \cdot \sigma_2$, where $\text{dom}(\sigma_1) = \{1, \dots, n\}$ and $\sigma_1(i) = \sigma(i)$, $i \in X(\sigma)$, $\sigma_1(i) = i$, $i \notin X(\sigma)$; $\text{dom}(\sigma_2) = \text{dom}(\sigma)$ and $\sigma_2(i) = i$, $i \in X(\sigma)$, $\sigma_2(i) = \sigma(i)$, $i \in \text{dom}(\sigma) \setminus X(\sigma)$. It follows immediately from the definition that $\sigma_1 \cdot \sigma_2 = \sigma_2 \cdot \sigma_1$. It is further easy to see (see e.g. [GK]) that

$$O(\sigma_1) \leq O(\sigma) \leq O(\sigma_1) + n - |X(\sigma)|, \quad O(\sigma_1) \leq \text{IO}(\sigma) \leq O(\sigma_1) + 2(n - |X(\sigma)|). \quad (7)$$

For a random element $\sigma \in \mathcal{IS}_n$, let $O'_n(\sigma) = O(\sigma_1)$. Given $X_n = X(\sigma) = k$, this has the same distribution as the order \tilde{O}_k of a random permutation of length k . In [ET] it was shown that, as $k \rightarrow \infty$,

$$\frac{\ln \tilde{O}_k - \frac{1}{2} \ln^2 k}{\sqrt{\frac{1}{3} \ln^3 k}} \xrightarrow{d} N(0, 1).$$

Hence, as $n \rightarrow \infty$,

$$\frac{\ln O'_n - \frac{1}{2} \ln^2 X_n}{\sqrt{\frac{1}{3} \ln^3 X_n}} \xrightarrow{d} N(0, 1),$$

and it follows as in the proof of Proposition 3 that

$$\frac{\ln O'_n - \frac{1}{8} \ln^2 n}{\sqrt{\frac{1}{24} \ln^3 n}} \xrightarrow{d} N(0, 1). \quad (8)$$

In particular, for almost all $\sigma \in \mathcal{IS}_n$ we have that $O(\sigma_1) \approx n^{(\ln n)/8}$. Since the difference between the left and the right sides of the inequalities in (7) is less than $2n$, in particular is $o(n^{(\ln n)/9})$, we obtain that, asymptotically, the left and the right sides of inequalities in (7) are the same. Now the necessary statement follows from (8). \square

6 Some generating functions

Consider some “objects” consisting of “components”, whose order in the objects is not important. Assume that there are a_m possible components containing exactly m elements. Let f_n denote the total number of objects, which consist of exactly n elements. The following well-known statement can be easily derived for example from [Wi, Theorem 3.4.1]

Proposition 6. *The exponential generating function for $\{f_n, n \geq 0\}$ is $F(z) = e^{A(z)}$, where $A(z) = \sum_{m=1}^{\infty} \frac{a_m}{m!} z^m$.*

Proposition 6 now can be used to compute the exponential generating functions for $|T_n|$, $|\mathcal{IS}_n|$.

Theorem 4. 1. *The exponential generating function for $a_n = |T_n|$ is $E_{T_n}(z) = e^{z/(1-z)}$.*

2. *The exponential generating function for $b_n = |\mathcal{IS}_n|$ is $E_{\mathcal{IS}_n}(z) = \frac{1}{1-z} e^{z/(1-z)}$.*

Proof. For T_n we have that components are chains and $a_m = m!$. Hence $A(z) = \sum_{m \geq 1} z^m = z/(1-z)$ and we get $F(z) = e^{z/(1-z)}$.

For \mathcal{IS}_n we have two types of components: cycles and chains, and thus $a_m = m! + (m-1)!$. This gives $A(z) = \frac{1}{1-z} - \ln(1-z)$ and therefore $F(z) = \frac{1}{1-z} e^{z/(1-z)}$. \square

Analogous arguments can be used to compute the exponential generating function for $L^{(n)}$ and L_n :

Theorem 5. 1. The exponential generating function for the sequence $c_n = |L^{(n)}|$ is $E_{L^{(n)}}(z) = \frac{z}{1-z} e^{z/(1-z)}$.

2. The exponential generating function for $d_n = |L_n|$ is $E_{L_n}(z) = \frac{z}{(1-z)^2} e^{z/(1-z)}$.

Proof. A fixed chain of length m is contained in exactly $|T_{n-m}|$ elements of T_n , and in exactly $|\mathcal{IS}_{n-m}|$ elements of \mathcal{IS}_n . This implies that $E_{L^{(n)}}(z) = \frac{z}{1-z} E_{T_n}(z) = \frac{z}{1-z} e^{z/(1-z)}$ and $E_{L_n}(z) = \frac{z}{1-z} E_{\mathcal{IS}_n}(z) = \frac{z}{(1-z)^2} e^{z/(1-z)}$. \square

Theorem 4 and the first part of Theorem 5 can now be used to derive the following corollaries:

Corollary 5. ([GM2, Theorem 7(2)]) $|\mathcal{IS}_n| = |T_n| + L^{(n)}$.

Proof. Follows from $E_{\mathcal{IS}_n}(z) = E_{T_n}(z) + E_{L^{(n)}}(z)$. \square

Corollary 6. ([GM2, Theorem 6(1)]) $|T_n| = \frac{1}{n} L_n$.

Proof. For the sequence $n|T_n|$ we have

$$E_{n|T_n|}(z) = z E'_{|T_n|}(z) = \frac{z}{(1-z)^2} e^{z/(1-z)} = E_{L_n}(z)$$

and the statement follows. \square

Corollary 7. ([GM2, Theorem 6(2)]) $|\mathcal{IS}_n| = \frac{1}{n+1} L^{(n+1)}$.

Proof. The statement is equivalent to $z E_{\mathcal{IS}_n}(z) = E_{L^{(n+1)}}(z)$, which is straightforward. \square

We also obtain one relation, which seems to be missing in [GM2].

Corollary 8. The total number P_n of fixed points for all elements from \mathcal{IS}_n satisfies $P_n = L^{(n)}$.

Proof. For $x \in \{1, \dots, n\}$, the point x is fixed in exactly $|\mathcal{IS}_{n-1}|$ elements of \mathcal{IS}_n , which implies that $E_{P_n}(z) = z E_{\mathcal{IS}_n}(z)$. Further $z E_{\mathcal{IS}_n}(z) = E_{L^{(n+1)}}(z)$ by Corollary 7 and the statement follows. \square

Acknowledgments

For the second author the research was partially supported by The Swedish Research Council and The Swedish Institute.

References

- [BH] A.D.Barbour, P.Hall, On the rate of Poisson convergence. Math. Proc. Cambridge Philos. Soc. 95 (1984), no. 3, 473–480.
- [BHJ] A.Barbour, L.Holst, S.Janson, Poisson approximation. Oxford Studies in Probability, 2. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1992. x+277 pp.
- [ET] P.Erdős, P.Turán, On some problems of a statistical group theory. I. Z. Wahrscheinlichkeitstheorie und Verw. Gebiete 4 1965 175–186 (1965).
- [GK] O.G.Ganyushkin, T.V.Kormysheva, Chain decomposition of partial permutations and classes of conjugated elements of the semigroup \mathcal{IS}_n , Proc. Kyiv University, Ser. Math.-Mech. 27 (1992), 10-18.
- [GM1] O.Ganyushkin, V.Mazorchuk, The full finite Inverse symmetric semigroup \mathcal{IS}_n , Preprint 2001:37, Chalmers University of Technology and Göteborg University, Göteborg, 2001.
- [GM2] O.Ganyushkin, V.Mazorchuk, Combinatorics and distributions of partial injections, Preprint 2003:14, Uppsala University, Sweden, 2003.
- [Go] V.L.Goncharov, Some facts from combinatorics, Izvestia Akad. Nauk. SSSR, Ser. Mat. 8: 3-48 (1944).
- [Li] S.Lipscomb, Symmetric inverse semigroups. Mathematical Surveys and Monographs, 46. American Mathematical Society, Providence, RI, 1996.
- [ST] B.M.Schein, B.Teclezghi, Endomorphisms of finite symmetric inverse semigroups. J. Algebra 198 (1997), no. 1, 300–310.
- [Wi] H.S.Wilf, Generatingfunctionology. Second edition. Academic Press, Inc., Boston, MA, 1994. x+228 pp.

S.J.: Department of Mathematics, Uppsala University, Box 480, SE 751 06, Uppsala, SWEDEN, e-mail: svante@math.uu.se, web: “<http://www.math.uu.se/~svante>”

V.M.: Department of Mathematics, Uppsala University, Box 480, SE 751 06, Uppsala, SWEDEN, e-mail: mazor@math.uu.se, web: “<http://www.math.uu.se/~mazor>”