Uppsala Universitet
Matematiska Institutionen
Thomas Erlandsson

DIFFERENTIAL GEOMETRY MN1 FALL 2001

PROBLEMS

9. Let $f : (a,b) \times (c,d) \to R^3$ be a surface in R^3 with constant Gauss curvature $K < 0$, defined in asymptotic coordinates, and parametrized by arc length so that

$$g_{11}(u) = g_{22}(u) = 1.$$

Let $\omega(u_0^1, u_0^2)$ be the unique number $0 < \omega(u_0^1, u_0^2) < \pi$ such that $\omega(u_0^1, u_0^2)$ is the angle between $f_{u^1}(u^1, u_0^2)|_{u^1=u_0^1}$ and $f_{u^2}(u_0^1, u^2)|_{u^2=u_0^2}$, i.e.

$$g_{12}(u) = \cos\omega(u).$$

a) Show that ω satisfies the differential equation

$$\frac{\partial^2\omega}{\partial u^1 \partial u^2} = (-K)\sin\omega$$

Hint: Use Gauss' equation which in the coordinates of the problem is

$$K = \frac{1}{2\sqrt{1-(g_{12})^2}}\left[\frac{\partial}{\partial u^1}\left(\frac{g_{12,2}}{\sqrt{1-(g_{12})^2}}\right) + \frac{\partial}{\partial u^2}\left(\frac{g_{12,1}}{\sqrt{1-(g_{12})^2}}\right)\right]$$

b) Show that every polygon Q with four sides and which is bounded by parameter curves has the area

$$\frac{1}{-K}\left(\sum_{i=1}^{4}\alpha_i - 2\pi\right) \leq \frac{2\pi}{-K},$$

where α_i are the inner angles in Q.

Hint: The area element is $\sqrt{g_{11}g_{22} - (g_{12})^2} = \sin\omega\, du^1 du^2$.

10. (For VG only)

The Poincaré Upper Half Plane Model of H^2

The surface H^2 is in the Poincaré upper half plane model the set

$$U = \{(u, v) \in R^2 \,|\, v > 0\}$$

with the Riemann metric

$$ds^2 = \frac{du^2 + dv^2}{v^2}.$$

Using Gauss' equation we find immediately that this surface has constant Gauss curvature $K = -1$. The line element ds^2 in H^2 is equal to the euclidean line element $du^2 + dv^2$ multiplied by a strictly positive function. Therefore an angle measured with respect to the Riemann metric coincides with the euclidean angle.

The geodesics in the upper half plane model of H^2 are the euclidean circles and straight lines which meet the boundary $v = 0$ orthogonally. This can be shown in the following way:

In H^2 we get $g_{11} = 1/v^2, \quad g_{12} = 0, \quad g_{22} = 1/v^2$ and it follows that

$$\Gamma^1_{11} = \Gamma^1_{22} = \Gamma^2_{12} = 0, \quad \Gamma^2_{11} = -\Gamma^2_{22} = -\Gamma^1_{21} = 1/v.$$

The differential equations of the geodesics can therefore be written

$$\ddot{u} - \frac{2\dot{u}\dot{v}}{v} = 0, \quad \ddot{v} + \frac{\dot{u}^2 - \dot{v}^2}{v} = 0.$$

If $\dot{u} = 0$ then $u =$ constant. In this case it is clear that the geodesic is a straight euclidean line orthogonal to $v = 0$.

If $\dot{u} \neq 0$ we get from the first equation that $\ln(\dot{u}/v^2) =$ constant so $\dot{u} = cv^2 \neq 0$ for some constant c. In the same way we get from the second equation that $\dot{u}^2 + \dot{v}^2 = bv^2 > 0$ for some constant b. By combining these equations we get $(dv/du)^2 = \dot{v}^2/\dot{u}^2 = b/c^2v^2 - 1$. Therefore $(u - a)^2 + v^2 = b/c^2$ for some constant a. This is a circle with centre on $v = 0$ and so meets $v = 0$ orthogonally.

The isometries of H^2 are well-known maps in the upper half plane model. Let $SL(2,R)$ be the special linear group in dimension 2, i.e. the group of all real (2×2)-matrices with determinant $=1$. $SL(2,R)$ acts on H^2 in the following way. Let $z=u+iv$. The points (u,v) in the upper half plane correspond to $z=u+iv,\ \ v>0$. If

$$g=\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2,R),$$

let

$$gz=\frac{az+b}{cz+d}.$$

Proposition

The group $SL(2,R)$ acts as a group of isometries on H^2.

Proof: Let $u+iv=z$ och

$$\frac{az+b}{cz+d}=\tilde{z}.$$

If we write $dz\,d\bar{z}$ for du^2+dv^2 the line element of H^2 can be written

$$ds^2(z)=\frac{-4dz\,d\bar{z}}{(z-\bar{z})^2}, \qquad \bar{z}=u-iv.$$

As $d\tilde{z}=d((az+b)(cz+d))=dz/((cz+d)^2$ it follows that $ds^2(z)=ds^2(\tilde{z})$, which means that $z\mapsto\tilde{z}$ is an isometry.

a) Calculate the arc length of the geodesic $c(t)=(r\cos t, r\sin t)$, $0<t<\pi$ starting from the top of the half circle , $t=\pi/2$. (**Result:** $|\ln\tan\frac{t}{2}|$)

Calculate also the arc length of the geodesic $u=u_0$ from $v=a$ till $v=b$. (**Result:** $(|\ln\frac{a}{b}|)$

b) Calculate the geodesic curvature k_g of the curve $v=1$. (**Result:** $k_g=1$)

The Poincaré Disc Model

The surface H^2 in the Poincaré disc model is the set

$$U = \{(u, v) \in R^2 | u^2 + v^2 < 4\},$$

with the Riemann metric

$$ds^2 = \left(1 - \frac{u^2 + v^2}{4}\right)^{-2} (du^2 + dv^2).$$

Using Gauss' equation we find that the surface has constant Gauss curvature $K = -1$. The geodesics in the disc model correspond to the circles orthogonal to the boundary of the disc and the diameters. This is most easily seen by showing that the map

$$w = \frac{z + 2i}{iz + 2}$$

is an isometry of the disc model onto the half plane model.

d) Calculate the arc length r of the geodesic $c(t) = (t\cos\vartheta, t\sin\vartheta)$, $0 \le t < 2$ beginning at origo.

(**Result:** $r = \ln\dfrac{2+t}{2-t} = 2\dfrac{1}{2}\ln\dfrac{1+t/2}{1-t/2} = 2\tanh^{-1}(t/2)$)

e) Show that in geodesic polar coordinates

$$ds^2 = dr^2 + \sinh^2(r)\, d\theta^2.$$

Hint: From the problem above follows that $t = 2\tanh(r/2)$.
Use $u = t\cos\theta, v = t\sin\theta$ and the given metric.